

This Page Is Inserted by IFW Operations  
and is not a part of the Official Record

## **BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,  
please do not report the images to the  
Image Problems Mailbox.**

This Page Blank (uspto)

DAN



GB00/3556.



PCT/GB 00/03556.

REC 2 OCT 2000	
WIPO	POSTOR IN PEOPLE

The Patent Office  
Concept House  
Cardiff Road  
Newport  
South Wales  
NP10 8QQ

10/088854


I, the undersigned, being an officer duly authorised in accordance with Section 74(1) and (4) of the Deregulation & Contracting Out Act 1994, to sign and issue certificates on behalf of the Comptroller-General, hereby certify that annexed hereto is a true copy of the documents as originally filed in connection with the patent application identified therein.

I also certify that by virtue of an assignment registered under the Patents Act 1977, the application is now proceeding in the name as substituted.

In accordance with the Patents (Companies Re-registration) Rules 1982, if a company named in this certificate and any accompanying documents has re-registered under the Companies Act 1980 with the same name as that with which it was registered immediately before re-registration save for the substitution as, or inclusion as, the last part of the name of the words "public limited company" or their equivalents in Welsh, references to the name of the company in this certificate and any accompanying documents shall be treated as references to the name with which it is so re-registered.

In accordance with the rules, the words "public limited company" may be replaced by p.l.c., plc, P.L.C. or PLC.

Re-registration under the Companies Act does not constitute a new legal entity but merely subjects the company to certain additional company law rules.



## PRIORITY DOCUMENT

SUBMITTED OR TRANSMITTED IN  
COMPLIANCE WITH RULE 17.1(a) OR (b)

Signed

Andrew Gersey

Dated

22 September 2000

---

This Page Blank (uspto)

GB9922156.6

---

By virtue of a direction given under Section of the Patents Act 1977, the application is proceeding in the name of

ASTRAZENECA AB,  
Incorporated in Sweden,  
S-151 85 Sodertalje,  
Sweden

[ADP No. 07822448003]

---

This Page Blank (uspto)

GB9922156.6

---

By virtue of a direction given under Section of the Patents Act 1977, the application is proceeding in the name of

ASTRAZENECA UK LIMITED  
Incorporated in the United Kingdom  
15 Stanhope Gate  
LONDON  
W1K 6LN  
United Kingdom

[ADP No. 07810294001]

---

This Page Blank (uspto)




Patents room THE PATENT OFFICE

Art 1077

21 SEP 1999

The  
Patent  
Office



# Request for grant of a patent

(See the notes on the back of this form. You can also get an explanatory leaflet from the Patent Office to help you fill in this form)

The Patent Office

Cardiff Road  
Newport  
Gwent NP9 1RH

1. Your reference

PHM 99-143

2. Patent application number

(The Patent Office will fill in this part)

9922156.6

21 SEP 1999

3. Full name, address and postcode of the or of each applicant (underline all surnames)

Zeneca Limited  
15 Stanhope Gate  
LONDON  
W1Y 6LW

SECTION 30 (1977 ACT) APPLICATION FILED 9/3/00

Patents ADP number (*if you know it*)

6254007002

If the applicant is a corporate body, give the country/state of its incorporation

4. Title of the invention

CHEMICAL COMPOUNDS

5. Name of your agent (*if you have one*)

BILL, Kevin

"Address for service" in the United Kingdom to which all correspondence should be sent (*including the postcode*)

AstraZeneca PLC  
Global Intellectual Property  
Mereside, Alderley Park,  
Macclesfield, Cheshire, SK10 4TG, GB

Patents ADP number (*if you know it*)

4469847002

6. If you are declaring priority from one or more earlier patent applications, give the country and the date of filing of the or of each of these earlier applications and (*if you know it*) the or each application number

Country

Priority application number  
(*if you know it*)

Date of filing  
(*day / month / year*)

7. If this application is divided or otherwise derived from an earlier UK application, give the number and the filing date of the earlier application

Number of earlier application

Date of filing  
(*day / month / year*)

8. Is a statement of inventorship and of right to grant of a patent required in support of this request? (*Answer 'Yes' if:*

- a) any applicant named in part 3 is not an inventor, or
- b) there is an inventor who is not named as an applicant, or

See notes (10/77)

**Patents Form 1/77**

Following marks must be made in the following boxes.  
Do not count copies of the same document

Continuation sheets of this form

Description

40

Claim(s)

Abstract

Drawing(s)

10. If you are also filing any of the following,  
state how many against each item.

Priority documents

Translations of priority documents

Statement of inventorship and right  
to grant of a patent (*Patents Form 7/77*)

Request for preliminary examination  
and search (*Patents Form 9/77*)

Request for substantive examination  
(*Patents Form 10/77*)

Any other documents  
(*please specify*)

11.

I/We request the grant of a patent on the basis of this application.

Signature

Date

Lynda May Slack 20 September 1999

12. Name and daytime telephone number of  
person to contact in the United Kingdom

Mrs Lynda May Slack 01625 516173

**Warning**

*After an application for a patent has been filed, the Comptroller of the Patent Office will consider whether publication or communication of the invention should be prohibited or restricted under Section 22 of the Patents Act 1977. You will be informed if it is necessary to prohibit or restrict your invention in this way. Furthermore, if you live in the United Kingdom, Section 23 of the Patents Act 1977 stops you from applying for a patent abroad without first getting written permission from the Patent Office unless an application has been filed at least 6 weeks beforehand in the United Kingdom for a patent for the same invention and either no direction prohibiting publication or communication has been given, or any such direction has been revoked.*

**Notes**

- If you need help to fill in this form or you have any questions, please contact the Patent Office on 0645 500505.*
- Write your answers in capital letters using black ink or you may type them.*
- If there is not enough space for all the relevant details on any part of this form, please continue on a separate sheet of paper and write "see continuation sheet" in the relevant part(s). Any continuation sheet should be attached to this form.*
- If you have answered 'Yes' Patents Form 7/77 will need to be filed.*
- Once you have filled in the form you must remember to sign and date it.*

CHEMICAL COMPOUNDS

The present invention relates to certain quinazoline derivatives for use in the treatment of certain diseases in particular to proliferative disease such as cancer and in the preparation of medicaments for use in the treatment of proliferative disease, to novel quinazoline compounds and to processes for their preparation, as well as pharmaceutical compositions containing them as active ingredient.

Cancer (and other hyperproliferative disease) is characterised by uncontrolled cellular proliferation. This loss of the normal regulation of cell proliferation often appears to occur as the result of genetic damage to cellular pathways that control progress through the cell cycle.

In eukaryotes, the cell cycle is largely controlled by an ordered cascade of protein phosphorylation. Several families of protein kinases that play critical roles in this cascade have now been identified. The activity of many of these kinases is increased in human tumours when compared to normal tissue. This can occur by either increased levels of expression of the protein (as a result of gene amplification for example), or by changes in expression of co activators or inhibitory proteins.

The first identified, and most widely studied of these cell cycle regulators have been the cyclin dependent kinases (or CDKs). Activity of specific CDKs at specific times is essential for both initiation and coordinated progress through the cell cycle. For example, the CDK4 protein appears to control entry into the cell cycle (the G0-G1-S transition) by phosphorylating the retinoblastoma gene product pRb. This stimulates the release of the transcription factor E2F from pRb, which then acts to increase the transcription of genes necessary for entry into S phase. The catalytic activity of CDK4 is stimulated by binding to a partner protein, Cyclin D. One of the first demonstrations of a direct link between cancer and the cell cycle was made with the observation that the Cyclin D1 gene was amplified and cyclin D protein levels increased (and hence the activity of CDK4 increased) in many human tumours (Reviewed in Sherr, 1996, Science 274: 1672-1677; Pines, 1995, Seminars in Cancer Biology 6: 63-72). Other studies (Loda et al., 1997, Nature Medicine 3(2): 231-234; Gemma et al., 1996, International Journal of Cancer 68(5): 605-11; Elledge et al. 1996, Trends in Cell Biology 6; 388-392) have shown that negative regulators of CDK function are frequently

down regulated or deleted in human tumours again leading to inappropriate activation of these kinases.

More recently, protein kinases that are structurally distinct from the CDK family have been identified which play critical roles in regulating the cell cycle and which also appear to be important in oncogenesis. These include the newly identified human homologues of the


*Drosophila* aurora and *S.cerevisiae* Ipl1 proteins. *Drosophila* aurora and *S.cerevisiae* Ipl1, which are highly homologous at the amino acid sequence level, encode serine/threonine protein kinases. Both aurora and Ipl1 are known to be involved in controlling the transition from the G2 phase of the cell cycle through mitosis, centrosome function, formation of a mitotic spindle and proper chromosome separation / segregation into daughter cells. The two human homologues of these genes, termed aurora1 and aurora2, encode cell cycle regulated protein kinases. These show a peak of expression and kinase activity at the G2/M boundary (aurora2) and in mitosis itself (aurora1). Several observations implicate the involvement of human aurora proteins, and particularly aurora2 in cancer. The aurora2 gene maps to chromosome 20q13, a region that is frequently amplified in human tumours including both breast and colon tumours. Aurora2 may be the major target gene of this amplicon, since aurora2 DNA is amplified and aurora2 mRNA overexpressed in greater than 50% of primary human colorectal cancers. In these tumours aurora2 protein levels appear greatly elevated compared to adjacent normal tissue. In addition, transfection of rodent fibroblasts with human aurora2 leads to transformation, conferring the ability to grow in soft agar and form tumours in nude mice (Bischoff et al., 1998, The EMBO Journal. 17(11): 3052-3065). Other work (Zhou et al., 1998, Nature Genetics. 20(2): 189-93) has shown that artificial overexpression of aurora2 leads to an increase in centrosome number and an increase in aneuploidy.

Importantly, it has also been demonstrated that abrogation of aurora2 expression and function by antisense oligonucleotide treatment of human tumour cell lines (WO 97/22702 and WO 99/37788) leads to cell cycle arrest in the G2 phase of the cell cycle and exerts an antiproliferative effect in these tumour cell lines. This indicates that inhibition of the function of aurora2 will have an antiproliferative effect that may be useful in the treatment of human tumours and other hyperproliferative diseases.

A number of quinazoline derivatives have been proposed hitherto for use in the inhibition of various kinases. For example, WO 96/09294, WO 96/15118 and WO 99/06378 describe the use of certain quinazoline compounds as receptor tyrosine kinase inhibitors, which may be useful in the treatment of proliferative disease.

The applicants have found a series of compounds which inhibit the effect of the aurora2 kinase and which are thus of use in the treatment of proliferative disease such as cancer, in particular in such diseases such as colorectal or breast where aurora 2 kinase is known to be active.

The present invention provides the use of a compound of formula (I)



(I)

or a salt, ester or amide thereof;

where X is O, or S, S(O) or S(O)<sub>2</sub>, NH or NR<sup>8</sup> where R<sup>8</sup> is hydrogen or C<sub>1-6</sub>alkyl;

Z is O or S,

n is 0, or an integer of from 1 to 6

R<sup>5</sup> is hydrogen or optionally substituted hydrocarbyl or optionally substituted heterocyclyl;

and R<sup>6</sup> and R<sup>7</sup> are independently selected from hydrogen, halo, C<sub>1-4</sub>alkyl, C<sub>1-4</sub>alkoxy, C<sub>1-</sub>

<sub>4</sub>alkoxymethyl, di(C<sub>1-4</sub>alkoxy)methyl, C<sub>1-4</sub>alkanoyl, trifluoromethyl, cyano, amino, C<sub>2-3</sub>alkenyl,

C<sub>2-3</sub>alkynyl, a phenyl group, a benzyl group or a 5-6-membered heterocyclic group with 1-3

heteroatoms, selected independently from O, S and N, which heterocyclic group may be

aromatic or non-aromatic and may be saturated (linked via a ring carbon or nitrogen atom) or

unsaturated (linked via a ring carbon atom), and which phenyl, benzyl or heterocyclic group

may bear on one or more ring carbon atoms up to 5 substituents selected from hydroxy, halogeno, C<sub>1-3</sub>alkyl, C<sub>1-3</sub>alkoxy, C<sub>1-3</sub>alkanoyloxy, trifluoromethyl, cyano, amino, nitro, C<sub>2-4</sub>alkanoyl, C<sub>1-4</sub>alkanoylamino, C<sub>1-4</sub>alkoxycarbonyl, C<sub>1-4</sub>alkylsulphanyl, C<sub>1-4</sub>alkylsulphinyl, C<sub>1-4</sub>alkylsulphonyl, carbamoyl, N-C<sub>1-4</sub>alkylcarbamoyl, N,N-di(C<sub>1-4</sub>alkyl)carbamoyl, aminosulphonyl, N-C<sub>1-4</sub>alkylaminosulphonyl, N,N-di(C<sub>1-4</sub>alkyl)aminosulphonyl, C<sub>1-4</sub>alkylsulphonylamino, and a saturated heterocyclic group selected from morpholino, thiomorpholino, pyrrolidinyl, piperazinyl, piperidinyl imidazolidinyl and pyrazolidinyl, which saturated heterocyclic group may bear 1 or 2 substituents selected from oxo, hydroxy, halogeno, C<sub>1-3</sub>alkyl, C<sub>1-3</sub>alkoxy, C<sub>1-3</sub>alkanoyloxy, trifluoromethyl, cyano, amino, nitro and C<sub>1-4</sub>alkoxycarbonyl, and

R<sup>1</sup>, R<sup>2</sup>, R<sup>3</sup>, R<sup>4</sup> are independently selected from, halo, cyano, nitro, trifluoromethyl, C<sub>1-3</sub>alkyl, -NR<sup>9</sup>R<sup>10</sup> (wherein R<sup>9</sup> and R<sup>10</sup>, which may be the same or different, each represents hydrogen or C<sub>1-3</sub>alkyl), or -X<sup>1</sup>R<sup>11</sup> (wherein X<sup>1</sup> represents a direct bond, -O-, -CH<sub>2</sub>-, -OCO-, carbonyl, -S-, -SO-, -SO<sub>2</sub>-, -NR<sup>12</sup>CO-, -CONR<sup>12</sup>-, -SO<sub>2</sub>NR<sup>12</sup>-, -NR<sup>13</sup>SO<sub>2</sub>- or -NR<sup>14</sup>- (wherein R<sup>12</sup>, R<sup>13</sup> and R<sup>14</sup> each independently represents hydrogen, C<sub>1-3</sub>alkyl or C<sub>1-3</sub>alkoxyC<sub>2-3</sub>alkyl), and R<sup>11</sup> is selected from one of the following eighteen groups:

1) hydrogen or C<sub>1-5</sub>alkyl which may be unsubstituted or which may be substituted with one or more groups selected from hydroxy, fluoro or amino,

2) C<sub>1-5</sub>alkylX<sup>2</sup>COR<sup>15</sup> (wherein X<sup>2</sup> represents -O- or -NR<sup>16</sup>- (in which R<sup>15</sup> represents hydrogen, C<sub>1-3</sub>alkyl or C<sub>1-3</sub>alkoxyC<sub>2-3</sub>alkyl) and R<sup>16</sup> represents C<sub>1-3</sub>alkyl, -NR<sup>17</sup>R<sup>18</sup> or -OR<sup>19</sup> (wherein R<sup>17</sup>, R<sup>18</sup> and R<sup>19</sup> which may be the same or different each represents hydrogen, C<sub>1-3</sub>alkyl or C<sub>1-3</sub>alkoxyC<sub>2-3</sub>alkyl));

3) C<sub>1-5</sub>alkylX<sup>3</sup>R<sup>20</sup> (wherein X<sup>3</sup> represents -O-, -S-, -SO-, -SO<sub>2</sub>-, -OCO-, -NR<sup>21</sup>CO-, -CONR<sup>22</sup>-, -SO<sub>2</sub>NR<sup>23</sup>-, -NR<sup>24</sup>SO<sub>2</sub>- or -NR<sup>25</sup>- (wherein R<sup>21</sup>, R<sup>22</sup>, R<sup>23</sup>, R<sup>24</sup> and R<sup>25</sup> each independently

represents hydrogen, C<sub>1-3</sub>alkyl or C<sub>1-3</sub>alkoxyC<sub>2-3</sub>alkyl) and R<sup>20</sup> represents hydrogen, C<sub>1-3</sub>alkyl, cyclopentyl, cyclohexyl or a 5-6-membered saturated heterocyclic group with 1-2 heteroatoms, selected independently from O, S and N, which C<sub>1-3</sub>alkyl group may bear 1 or 2 substituents selected from oxo, hydroxy, halogeno and C<sub>1-4</sub>alkoxy and which cyclic group may bear 1 or 2 substituents selected from oxo, hydroxy, halogeno, C<sub>1-4</sub>alkyl, C<sub>1-4</sub>hydroxyalkyl and C<sub>1-4</sub>alkoxy);

- 4)  $C_{1-5}alkylX^4C_{1-5}alkylX^5R^{26}$  (wherein  $X^4$  and  $X^5$  which may be the same or different are each -O-, -S-, -SO-, -SO<sub>2</sub>-, -NR<sup>27</sup>CO-, -CONR<sup>28</sup>-, -SO<sub>2</sub>NR<sup>29</sup>-, -NR<sup>30</sup>SO<sub>2</sub>- or -NR<sup>31</sup>- (wherein R<sup>27</sup>, R<sup>28</sup>, R<sup>29</sup>, R<sup>30</sup> and R<sup>31</sup> each independently represents hydrogen, C<sub>1-3</sub>alkyl or C<sub>1-3</sub>alkoxyC<sub>2-3</sub>alkyl) and R<sup>26</sup> represents hydrogen or C<sub>1-3</sub>alkyl);
- 5) R<sup>32</sup> (wherein R<sup>32</sup> is a 5-6-membered saturated heterocyclic group (linked via carbon or nitrogen) with 1-2 heteroatoms, selected independently from O, S and N, which heterocyclic group may bear 1 or 2 substituents selected from oxo, hydroxy, halogeno, C<sub>1-4</sub>alkyl, C<sub>1-4</sub>hydroxyalkyl, C<sub>1-4</sub>alkoxy, C<sub>1-4</sub>alkoxyC<sub>1-4</sub>alkyl and C<sub>1-4</sub>alkylsulphonylC<sub>1-4</sub>alkyl);
- 6) C<sub>1-5</sub>alkylR<sup>32</sup> (wherein R<sup>32</sup> is as defined hereinbefore);
- 7) C<sub>2-5</sub>alkenylR<sup>32</sup> (wherein R<sup>32</sup> is as defined hereinbefore);
- 8) C<sub>2-5</sub>alkynylR<sup>32</sup> (wherein R<sup>32</sup> is as defined hereinbefore);
- 9) R<sup>33</sup> (wherein R<sup>33</sup> represents a pyridone group, a phenyl group or a 5-6-membered aromatic heterocyclic group (linked via carbon or nitrogen) with 1-3 heteroatoms selected from O, N and S, which pyridone, phenyl or aromatic heterocyclic group may carry up to 5 substituents on an available carbon atom selected from hydroxy, halogeno, amino, C<sub>1-4</sub>alkyl, C<sub>1-4</sub>alkoxy, C<sub>1-4</sub>hydroxyalkyl, C<sub>1-4</sub>aminoalkyl, C<sub>1-4</sub>alkylamino, C<sub>1-4</sub>hydroxyalkoxy, carboxy, trifluoromethyl, cyano, -CONR<sup>34</sup>R<sup>35</sup> and -NR<sup>36</sup>COR<sup>37</sup> (wherein R<sup>34</sup>, R<sup>35</sup>, R<sup>36</sup> and R<sup>37</sup>, which may be the same or different, each represents hydrogen, C<sub>1-4</sub>alkyl or C<sub>1-3</sub>alkoxyC<sub>2-3</sub>alkyl));
- 10) C<sub>1-5</sub>alkylR<sup>33</sup> (wherein R<sup>33</sup> is as defined hereinbefore);
- 11) C<sub>2-5</sub>alkenylR<sup>33</sup> (wherein R<sup>33</sup> is as defined hereinbefore);
- 12) C<sub>2-5</sub>alkynylR<sup>33</sup> (wherein R<sup>33</sup> is as defined hereinbefore);
- 13) C<sub>1-5</sub>alkylX<sup>6</sup>R<sup>33</sup> (wherein X<sup>6</sup> represents -O-, -S-, -SO-, -SO<sub>2</sub>-, -NR<sup>38</sup>CO-, -CONR<sup>39</sup>-, -SO<sub>2</sub>NR<sup>40</sup>-, -NR<sup>41</sup>SO<sub>2</sub>- or -NR<sup>42</sup>- (wherein R<sup>38</sup>, R<sup>39</sup>, R<sup>40</sup>, R<sup>41</sup> and R<sup>42</sup> each independently represents hydrogen, C<sub>1-3</sub>alkyl or C<sub>1-3</sub>alkoxyC<sub>2-3</sub>alkyl) and R<sup>33</sup> is as defined hereinbefore);
- 14) C<sub>2-5</sub>alkenylX<sup>7</sup>R<sup>33</sup> (wherein X<sup>7</sup> represents -O-, -S-, -SO-, -SO<sub>2</sub>-, -NR<sup>43</sup>CO-, -CONR<sup>44</sup>-, -SO<sub>2</sub>NR<sup>45</sup>-, -NR<sup>46</sup>SO<sub>2</sub>- or -NR<sup>47</sup>- (wherein R<sup>43</sup>, R<sup>44</sup>, R<sup>45</sup>, R<sup>46</sup> and R<sup>47</sup> each independently represents hydrogen, C<sub>1-3</sub>alkyl or C<sub>1-3</sub>alkoxyC<sub>2-3</sub>alkyl) and R<sup>33</sup> is as defined hereinbefore);
- 15) C<sub>2-5</sub>alkynylX<sup>8</sup>R<sup>33</sup> (wherein X<sup>8</sup> represents -O-, -S-, -SO-, -SO<sub>2</sub>-, -NR<sup>48</sup>CO-, -CONR<sup>49</sup>-, -SO<sub>2</sub>NR<sup>50</sup>-, -NR<sup>51</sup>SO<sub>2</sub>- or -NR<sup>52</sup>- (wherein R<sup>48</sup>, R<sup>49</sup>, R<sup>50</sup>, R<sup>51</sup> and R<sup>52</sup> each independently represents hydrogen, C<sub>1-3</sub>alkyl or C<sub>1-3</sub>alkoxyC<sub>2-3</sub>alkyl) and R<sup>33</sup> is as defined hereinbefore);

16)  $C_{1-3}alkylX^9C_{1-3}alkylR^{33}$  (wherein  $X^9$  represents -O-, -S-, -SO-, -SO<sub>2</sub>-, -NR<sup>53</sup>CO-, -CONR<sup>54</sup>-, -SO<sub>2</sub>NR<sup>55</sup>-, -NR<sup>56</sup>SO<sub>2</sub>- or -NR<sup>57</sup>- (wherein R<sup>53</sup>, R<sup>54</sup>, R<sup>55</sup>, R<sup>56</sup> and R<sup>57</sup> each independently represents hydrogen, C<sub>1-3</sub>alkyl or C<sub>1-3</sub>alkoxyC<sub>2-3</sub>alkyl) and R<sup>33</sup> is as defined hereinbefore); and  
 17)  $C_{1-3}alkylX^9C_{1-3}alkylR^{32}$  (wherein  $X^9$  and R<sup>32</sup> are as defined hereinbefore);

5 and R<sup>1</sup> and R<sup>4</sup> may additionally be hydrogen; in the preparation of a medicament for use in the inhibition of aurora 2 kinase. In particular, such medicaments are useful in the treatment of proliferative disease such as cancer, and in particular cancers where aurora 2 is upregulated such as colon or breast cancers.

In this specification the term 'alkyl' when used either alone or as a suffix includes  
 10 straight chained, branched structures. Unless otherwise stated, these groups may contain up to 10, preferably up to 6 and more preferably up to 4 carbon atoms. Similarly the terms "alkenyl" and "alkynyl" refer to unsaturated straight or branched structures containing for example from 2 to 10, preferably from 2 to 6 carbon atoms. Cyclic moieties such as cycloalkyl, cycloalkenyl and cycloalkynyl are similar in nature but have at least 3 carbon  
 15 atoms. Terms such as "alkoxy" comprise alkyl groups as is understood in the art.

The term "halo" includes fluoro, chloro, bromo and iodo. References to aryl groups include aromatic carbocyclic groups such as phenyl and naphthyl. The term "heterocyclyl" includes aromatic or non-aromatic rings, for example containing from 4 to 20, suitably from 5 to 8 ring atoms, at least one of which is a heteroatom such as oxygen, sulphur or nitrogen.

20 Examples of such groups include furyl, thienyl, pyrrolyl, pyrrolidinyl, imidazolyl, triazolyl, thiazolyl, tetrazolyl, oxazolyl, isoxazolyl, pyrazolyl, pyridyl, pyrimidinyl, pyrazinyl, pyridazinyl, triazinyl, quinolinyl, isoquinolinyl, quinoxalinyl, benzothiazolyl, benzoxazolyl, benzothienyl or benzofuryl.

"Heteroaryl" refers to those groups described above which have an aromatic character.  
 25 The term "aralkyl" refers to aryl substituted alkyl groups such as benzyl.

Other expressions used in the specification include "hydrocarbyl" which refers to any structure comprising carbon and hydrogen atoms. For example, these may be alkyl, alkenyl, alkynyl, aryl, heterocyclyl, alkoxy, aralkyl, cycloalkyl, cycloalkenyl or cycloalkynyl.

The term "functional group" refers to reactive substituents such as nitro, cyano, halo,  
 30 oxo, =CR<sup>78</sup>R<sup>79</sup>, C(O)<sub>x</sub>R<sup>77</sup>, OR<sup>77</sup>, S(O)<sub>y</sub>R<sup>77</sup>, NR<sup>78</sup>R<sup>79</sup>, C(O)NR<sup>78</sup>R<sup>79</sup>, OC(O)NR<sup>78</sup>R<sup>79</sup>, =NOR<sup>77</sup>, -NR<sup>77</sup>C(O)<sub>x</sub>R<sup>78</sup>, -NR<sup>77</sup>CONR<sup>78</sup>R<sup>79</sup>, -N=CR<sup>78</sup>R<sup>79</sup>, S(O)<sub>y</sub>NR<sup>78</sup>R<sup>79</sup> or -NR<sup>77</sup>S(O)<sub>y</sub>R<sup>78</sup> where R<sup>77</sup>,



$R^{78}$  and  $R^{79}$  are independently selected from hydrogen or optionally substituted hydrocarbyl, or  $R^{78}$  and  $R^{79}$  together form an optionally substituted ring which optionally contains further heteroatoms such as  $S(O)_y$ , oxygen and nitrogen,  $x$  is an integer of 1 or 2,  $y$  is 0 or an integer of 1-3.

5            Suitable optional substituents for hydrocarbyl groups  $R^{77}$ ,  $R^{78}$  and  $R^{79}$  include halo, perhaloalkyl such as trifluoromethyl, mercapto, hydroxy, carboxy, alkoxy, heteroaryl, heteroaryloxy, alkenyloxy, alkynyloxy, alkoxyalkoxy, aryloxy (where the aryl group may be substituted by halo, nitro, or hydroxy), cyano, nitro, amino, mono- or di-alkyl amino, oximino or  $S(O)_y$ , where  $y$  is as defined above.

10            Preferably  $R^1$  and  $R^4$  are hydrogen.

In a preferred embodiment, at least one group  $R^2$  or  $R^3$ , preferably  $R^3$ , comprises a chain of at least 3 and preferably at least 4 optionally substituted carbon atoms or heteroatoms such as oxygen, nitrogen or sulphur. Most preferably the chain is substituted by a polar group which assists in solubility.

15            Suitably  $R^3$  is a group  $XR^{11}$ . Preferably in this case,  $X^1$  is oxygen and  $R^{11}$  is selected from a group of formula (1) or (10) above. Particular groups  $R^{11}$  are those in group (1) above, especially alkyl such as methyl or halo substituted alkyl, or those in group (10) above. In one preferred embodiment, at least one of  $R^2$  or  $R^3$  is a group  $OC_{1-3}alkylR^{33}$  and  $R^{33}$  is a heterocyclic ring such as an N-linked morpholine ring, in particular 3-morpholinopropoxy.

20            Suitably  $R^2$  is selected from, halo, cyano, nitro, trifluoromethyl,  $C_{1-3}alkyl$ ,  $-NR^9R^{10}$  (wherein  $R^9$  and  $R^{10}$ , which may be the same or different, each represents hydrogen or  $C_{1-3}alkyl$ ), or a group  $-X^1R^{11}$ . Preferred examples of  $-X^1R^{11}$  for  $R^2$  include those listed above in relation to  $R^3$ .

Other examples for  $R^2$  and  $R^3$  include methoxy or 3,3,3-trifluoroethoxy.

25            Preferably  $X$  is NH or O and is most preferably NH.

Suitably  $R^5$  is hydrogen, optionally substituted alkenyl, optionally substituted aryl or optionally substituted heterocyclyl. In particular,  $R^5$  is hydrogen, ethenyl, optionally substituted phenyl or optionally substituted pyridyl. Suitable optional substituents  $R^5$  groups include  $C_{1-3}alkoxy$  such as methoxy,  $C_{1-3}alkyl$  such as methyl

30            Preferably  $n$  is 0 when  $R^5$  is optionally substituted phenyl or naphthyl. When  $R^5$  is hydrogen,  $n$  is suitably other than 0 and preferably from 1 to 5, more preferably from 3 to 5.

Suitably  $R^6$  and  $R^7$  are independently selected from hydrogen halo,  $C_{1-4}$ alkoxy such as methoxy, or ethoxy, cyano, trifluoromethyl, or phenyl.

Preferably  $R^6$  and  $R^7$  are hydrogen.

Suitable pharmaceutically acceptable salts of compounds of formula (I) include acid  
5 addition salts such as methanesulfonate, fumarate, hydrochloride, hydrobromide, citrate, maleate and salts formed with phosphoric and sulphuric acid. There may be more than one cation or anion depending on the number of charged functions and the valency of the cations or anions. Where the compound of formula (I) includes an acid functionality, salts may be base salts such as an alkali metal salt for example sodium, an alkaline earth metal salt for  
10 example calcium or magnesium, an organic amine salt for example triethylamine, morpholine, *N*-methylpiperidine, *N*-ethylpiperidine, procaine, dibenzylamine, *N,N*-dibenzylethylamine or amino acids for example lysine. A preferred pharmaceutically acceptable salt is a sodium salt.

An *in vivo* hydrolysable ester of a compound of the formula (I) containing carboxy or hydroxy group is, for example, a pharmaceutically acceptable ester which is hydrolysed in the  
15 human or animal body to produce the parent acid or alcohol.

Suitable pharmaceutically acceptable esters for carboxy include  $C_{1-6}$ alkyl esters such as methyl or ethyl esters,  $C_{1-6}$ alkoxymethyl esters for example methoxymethyl,  $C_{1-6}$ alkanoyloxymethyl esters for example pivaloyloxymethyl, phthalidyl esters,  $C_{3-8}$ cycloalkoxy-carbonyloxy $C_{1-6}$ alkyl esters for example 1-cyclohexylcarbonyloxyethyl;  
20 1,3-dioxolen-2-onylmethyl esters for example 5-methyl-1,3-dioxolen-2-onylmethyl; and  $C_{1-6}$ alkoxycarbonyloxyethyl esters for example 1-methoxycarbonyloxyethyl and may be formed at any carboxy group in the compounds of this invention.

An *in vivo* hydrolysable ester of a compound of the formula (I) containing a hydroxy group includes inorganic esters such as phosphate esters and  $\alpha$ -acyloxyalkyl ethers and related  
25 compounds which as a result of the *in vivo* hydrolysis of the ester breakdown to give the parent hydroxy group. Examples of  $\alpha$ -acyloxyalkyl ethers include acetoxymethoxy and 2,2-dimethylpropionyloxymethoxy. A selection of *in vivo* hydrolysable ester forming groups for hydroxy include alkanoyl, benzoyl, phenylacetyl and substituted benzoyl and phenylacetyl, alkoxycarbonyl (to give alkyl carbonate esters), dialkylcarbamoyl and  
30 *N*-(dialkylaminoethyl)-*N*-alkylcarbamoyl (to give carbamates), dialkylaminoacetyl and carboxyacetyl.

Suitable amides are derived from compounds of formula (I) which have a carboxy group which is derivatised into an amide such as a N-C<sub>1-6</sub>alkyl and N,N-di-(C<sub>1-6</sub>alkyl)amide such as N-methyl, N-ethyl, N-propyl, N,N-dimethyl, N-ethyl-N-methyl or N,N-diethylamide.

5 Esters which are not *in vivo* hydrolysable may be useful as intermediates in the production of the compounds of formula (I).

Particular examples of compounds of formula (I) are set out in Table 1































Table 1

Comp No	R <sup>2</sup>	R <sup>3</sup>	R <sup>5</sup>	R <sup>6</sup>	R <sup>7</sup>	n	Z	X
1	OCH <sub>3</sub>	OCH <sub>3</sub>	H	H	H	4	O	NH
2	OCH <sub>3</sub>	OCH <sub>3</sub>		CH <sub>3</sub>	H	1	O	NH
3	OCH <sub>3</sub>	OCH <sub>3</sub>		CH <sub>3</sub>	H	1	O	NH
4	OCH <sub>3</sub>	OCH <sub>3</sub>		CH <sub>3</sub>	H	1	O	NH
5	OCH <sub>3</sub>	OCH <sub>3</sub>		CH <sub>3</sub>	H	1	O	NH
6	OCOCH <sub>3</sub>	OCH <sub>3</sub>		F	H	1	O	NH
7	OCH <sub>3</sub>	OCH <sub>3</sub>	H	H	H	3	O	O

8	OCH <sub>3</sub>	OCH <sub>3</sub>		H	H	0	O	O
9	OCH <sub>3</sub>	OCH <sub>3</sub>		H	H	1	O	O
10	OCH <sub>3</sub>	OCH <sub>3</sub>	H	H	H	1	S	O
11	OCH <sub>3</sub>	OCH <sub>3</sub>	H	H	H	5	O	O
12	OCH <sub>3</sub>	OCH <sub>3</sub>		H	H	1	O	NH
13	OCH <sub>3</sub>	OCH <sub>3</sub>		H	H	2	O	NH
14	OCH <sub>3</sub>	OCH <sub>3</sub>	CH=CH <sub>2</sub>	H	H	1	O	NH
15	OCH <sub>3</sub>	O(CH <sub>2</sub> ) <sub>3</sub> N 		Cl	H	0	O	NH
16	OCH <sub>3</sub>	O(CH <sub>2</sub> ) <sub>3</sub> N 		Cl	H	0	S	NH
17	OCH <sub>3</sub>	O(CH <sub>2</sub> ) <sub>3</sub> N 	H	H	H	1	S	ONH
18	OCH <sub>3</sub>	O(CH <sub>2</sub> ) <sub>3</sub> N 		H	H	0	O	NH
19	OCH <sub>3</sub>	OCH <sub>2</sub> C <sub>6</sub> H <sub>5</sub>		H	H	0	O	NH
20	OCH <sub>3</sub>	O(CH <sub>2</sub> ) <sub>3</sub> N 		H	H	1	O	NH
21	OCH <sub>3</sub>	O(CH <sub>2</sub> ) <sub>3</sub> N 		H	H	0	S	NH
22	OCH <sub>3</sub>	O(CH <sub>2</sub> ) <sub>3</sub> N 	H	H	H	4	O	NH

23	OCH <sub>3</sub>	O(CH <sub>2</sub> ) <sub>3</sub> N 		H	H	0	O	NH
24	OCH <sub>3</sub>	OCH <sub>2</sub> CF <sub>3</sub>		F	H	1	O	NH
25	OCH <sub>3</sub>	OCH <sub>3</sub>		H	H	1	O	NH

Certain compounds of formula (I) are novel and these form a further aspect of the invention. Particular examples of such compounds are compounds of formula (IA)



(IA)

where X, Z, R<sup>5</sup>, R<sup>6</sup> and R<sup>7</sup> and n are as defined in relation to formula (I) and R<sup>66</sup> is halo, cyano, nitro, trifluoromethyl, C<sub>1-3</sub>alkyl, -NR<sup>9</sup>R<sup>10</sup> (wherein R<sup>9</sup> and R<sup>10</sup>, which may be the same or different, each represents hydrogen or C<sub>1-3</sub>alkyl), or a group -X<sup>1</sup>R<sup>11</sup> where X<sup>1</sup> and R<sup>11</sup> are as defined in relation to formula (I) and R<sup>11</sup> is particularly a group of sub group (1) or (10), and R<sup>67</sup> is C<sub>1-6</sub>alkoxy optionally substituted by fluorine or a group X<sup>1</sup>R<sup>33</sup> in which X<sup>1</sup> and R<sup>33</sup> are as defined in relation to formula (I), and in particular X<sup>1</sup> is oxygen and R<sup>33</sup> is or a 5-6-membered aromatic heterocyclic group (linked via nitrogen) with 1-3 heteroatoms selected from O, N and S; provided that at least one of R<sup>66</sup> and R<sup>67</sup> is other than unsubstituted methoxy.

A preferred example of R<sup>67</sup> is 3-morpholinopropoxy.

Preferably  $X^1$  is oxygen.


Preferably at least  $R^{67}$  is other than unsubstituted alkoxy.

Where  $R^{66}$  or  $R^{67}$  is unsubstituted alkoxy, it is preferably methoxy.

Suitable halo substituents for  $R^{66}$  and  $R^{67}$  are fluoro.


5 Other examples for  $R^{66}$  and/or  $R^{67}$  include 3,3,3-trifluoroethoxy.

Compounds of formula (I) may be prepared by methods known in the art or by analogous methods. For example, a compound of formula (I) can be prepared by reacting a compound of formula (VII)



(VII)

10 where  $R^{1'}$ ,  $R^{2''}$ ,  $R^{3''}$ , and  $R^{4'}$  are equivalent to a group  $R^1$ ,  $R^2$ ,  $R^3$  and  $R^4$  as defined in relation to formula (I) or a precursor thereof, and  $R^{85}$  is a leaving group, with a compound of formula (VIII)



(VIII)

15 where X, Z,  $R^5$ ,  $R^6$ ,  $R^7$  and n are as defined in relation to formula (I), and thereafter if desired or necessary converting a group  $R^{1'}$ ,  $R^{2''}$ ,  $R^{3''}$  or  $R^{4'}$  to a group  $R^1$ ,  $R^2$ ,  $R^3$  and  $R^4$  respectively or to a different such group.

20 Suitable leaving groups for  $R^{85}$  include halo such as chloro, mesylate and tosylate.

The reaction is suitably effected in an organic solvent such as an alcohol like isopropanol, at elevated temperatures, conveniently at the reflux temperature of the solvent.

The conversion of a group  $R^1$ ,  $R^2$ ,  $R^3$  or  $R^4$  to a group  $R^1$ ,  $R^2$ ,  $R^3$  and  $R^4$  respectively or to a different such group, may be particularly useful in connection with the preparation of compounds of formula (IA) and examples of these preparations are provided hereinafter.

Compounds of formula (VII) and (VIII) are either known compounds or they can be  
5 derived from known compounds by conventional methods.

---

Compounds of formula (I) are inhibitors of aurora 2 kinase. As a result, these compounds can be used to treat disease mediated by these agents, in particular proliferative disease.

According to a further aspect of the present invention there is provided a method for  
10 inhibiting aurora 2 kinase in a warm blooded animal, such as man, in need of such treatment, which comprises administering to said animal an effective amount of a compound of formula (I), or a pharmaceutically acceptable salt, or an *in vivo* hydrolysable ester thereof.

Novel compounds of formula (I) have not hitherto been proposed for use in therapy. Thus, according to a further aspect of the invention there is provided a compound of the  
15 formula (IA) as defined herein, or a pharmaceutically acceptable salt or an *in vivo* hydrolysable ester thereof, for use in a method of treatment of the human or animal body by therapy. In particular, the compounds are used in methods of treatment of proliferative disease such as cancer and in particular cancers such as colorectal or breast cancer where aurora 2 is upregulated.

20 Compounds of formula (I) are suitably applied in the form of a pharmaceutical composition. Preferred compounds of formula (I) for use in the compositions of the invention are as described above.

Some of these are novel and form yet a further aspect of the invention. Thus, the invention also provides a pharmaceutical composition comprising a compound of formula  
25 (IA) as defined herein, or a pharmaceutically acceptable salt, or an *in vivo* hydrolysable ester thereof, in combination with at pharmaceutically acceptable carrier.

The compositions of compounds of formula (I) may be in a form suitable for oral use (for example as tablets, lozenges, hard or soft capsules, aqueous or oily suspensions, emulsions, dispersible powders or granules, syrups or elixirs), for topical use (for example as  
30 creams, ointments, gels, or aqueous or oily solutions or suspensions), for administration by inhalation (for example as a finely divided powder or a liquid aerosol), for administration by

insufflation (for example as a finely divided powder) or for parenteral administration (for example as a sterile aqueous or oily solution for intravenous, subcutaneous, intramuscular or intramuscular dosing or as a suppository for rectal dosing).

The compositions of the invention may be obtained by conventional procedures using conventional pharmaceutical excipients, well known in the art. Thus, compositions intended for oral use may contain, for example, one or more colouring, sweetening, flavouring and/or preservative agents.

Suitable pharmaceutically acceptable excipients for a tablet formulation include, for example, inert diluents such as lactose, sodium carbonate, calcium phosphate or calcium carbonate, granulating and disintegrating agents such as corn starch or algenic acid; binding agents such as starch; lubricating agents such as magnesium stearate, stearic acid or talc; preservative agents such as ethyl or propyl p-hydroxybenzoate, and anti-oxidants, such as ascorbic acid. Tablet formulations may be uncoated or coated either to modify their disintegration and the subsequent absorption of the active ingredient within the gastrointestinal track, or to improve their stability and/or appearance, in either case, using conventional coating agents and procedures well known in the art.

Compositions for oral use may be in the form of hard gelatin capsules in which the active ingredient is mixed with an inert solid diluent, for example, calcium carbonate, calcium phosphate or kaolin, or as soft gelatin capsules in which the active ingredient is mixed with water or an oil such as peanut oil, liquid paraffin, or olive oil.

Aqueous suspensions generally contain the active ingredient in finely powdered form together with one or more suspending agents, such as sodium carboxymethylcellulose, methylcellulose, hydroxypropylmethylcellulose, sodium alginate, polyvinyl-pyrrolidone, gum tragacanth and gum acacia; dispersing or wetting agents such as lecithin or condensation products of an alkylene oxide with fatty acids (for example polyoxyethylene stearate), or condensation products of ethylene oxide with long chain aliphatic alcohols, for example heptadecaethyleneoxycetanol, or condensation products of ethylene oxide with partial esters derived from fatty acids and a hexitol such as polyoxyethylene sorbitol monooleate, or condensation products of ethylene oxide with long chain aliphatic alcohols, for example heptadecaethyleneoxycetanol, or condensation products of ethylene oxide with partial esters derived from fatty acids and a hexitol such as polyoxyethylene sorbitol monooleate, or



condensation products of ethylene oxide with partial esters derived from fatty acids and hexitol anhydrides, for example polyethylene sorbitan monooleate. The aqueous suspensions may also contain one or more preservatives (such as ethyl or propyl p-hydroxybenzoate, anti-oxidants (such as ascorbic acid), colouring agents, flavouring agents, and/or sweetening agents (such as sucrose, saccharine or aspartame).

---

Oily suspensions may be formulated by suspending the active ingredient in a vegetable oil (such as arachis oil, olive oil, sesame oil or coconut oil) or in a mineral oil (such as liquid paraffin). The oily suspensions may also contain a thickening agent such as beeswax, hard paraffin or cetyl alcohol. Sweetening agents such as those set out above, and flavouring agents may be added to provide a palatable oral preparation. These compositions may be preserved by the addition of an anti-oxidant such as ascorbic acid.

Dispersible powders and granules suitable for preparation of an aqueous suspension by the addition of water generally contain the active ingredient together with a dispersing or wetting agent, suspending agent and one or more preservatives. Suitable dispersing or wetting agents and suspending agents are exemplified by those already mentioned above. Additional excipients such as sweetening, flavouring and colouring agents, may also be present.

The pharmaceutical compositions of the invention may also be in the form of oil-in-water emulsions. The oily phase may be a vegetable oil, such as olive oil or arachis oil, or a mineral oil, such as for example liquid paraffin or a mixture of any of these. Suitable emulsifying agents may be, for example, naturally-occurring gums such as gum acacia or gum tragacanth, naturally-occurring phosphatides such as soya bean, lecithin, an esters or partial esters derived from fatty acids and hexitol anhydrides (for example sorbitan monooleate) and condensation products of the said partial esters with ethylene oxide such as polyoxyethylene sorbitan monooleate. The emulsions may also contain sweetening, flavouring and preservative agents.

Syrups and elixirs may be formulated with sweetening agents such as glycerol, propylene glycol, sorbitol, aspartame or sucrose, and may also contain a demulcent, preservative, flavouring and/or colouring agent.

The pharmaceutical compositions may also be in the form of a sterile injectable aqueous or oily suspension, which may be formulated according to known procedures using one or more of the appropriate dispersing or wetting agents and suspending agents, which

have been mentioned above. A sterile injectable preparation may also be a sterile injectable solution or suspension in a non-toxic parenterally-acceptable diluent or solvent, for example a solution in 1,3-butanediol.

Suppository formulations may be prepared by mixing the active ingredient with a suitable non-irritating excipient which is solid at ordinary temperatures but liquid at the rectal temperature and will therefore melt in the rectum to release the drug. Suitable excipients include, for example, cocoa butter and polyethylene glycols.

Topical formulations, such as creams, ointments, gels and aqueous or oily solutions or suspensions, may generally be obtained by formulating an active ingredient with a conventional, topically acceptable, vehicle or diluent using conventional procedure well known in the art.

Compositions for administration by insufflation may be in the form of a finely divided powder containing particles of average diameter of, for example,  $30\mu$  or much less, the powder itself comprising either active ingredient alone or diluted with one or more physiologically acceptable carriers such as lactose. The powder for insufflation is then conveniently retained in a capsule containing, for example, 1 to 50mg of active ingredient for use with a turbo-inhaler device, such as is used for insufflation of the known agent sodium cromoglycate.

Compositions for administration by inhalation may be in the form of a conventional pressurised aerosol arranged to dispense the active ingredient either as an aerosol containing finely divided solid or liquid droplets. Conventional aerosol propellants such as volatile fluorinated hydrocarbons or hydrocarbons may be used and the aerosol device is conveniently arranged to dispense a metered quantity of active ingredient.

For further information on Formulation the reader is referred to Chapter 25.2 in Volume 5 of Comprehensive Medicinal Chemistry (Corwin Hansch; Chairman of Editorial Board), Pergamon Press 1990.

The amount of active ingredient that is combined with one or more excipients to produce a single dosage form will necessarily vary depending upon the host treated and the particular route of administration. For example, a formulation intended for oral administration to humans will generally contain, for example, from 0.5 mg to 2 g of active agent compounded with an appropriate and convenient amount of excipients which may vary from

about 5 to about 98 percent by weight of the total composition. Dosage unit forms will generally contain about 1 mg to about 500 mg of an active ingredient. For further information on Routes of Administration and Dosage Regimes the reader is referred to Chapter 25.3 in Volume 5 of Comprehensive Medicinal Chemistry (Corwin Hansch; Chairman of Editorial Board), Pergamon Press 1990.

The size of the dose for therapeutic or prophylactic purposes of a compound of the Formula I will naturally vary according to the nature and severity of the conditions, the age and sex of the animal or patient and the route of administration, according to well known principles of medicine. As mentioned above, compounds of the Formula I are useful in treating diseases or medical conditions which are due alone or in part to the effects of aurora 2 kinase.

In using a compound of the Formula I for therapeutic or prophylactic purposes it will generally be administered so that a daily dose in the range, for example, 0.5 mg to 75 mg per kg body weight is received, given if required in divided doses. In general lower doses will be administered when a parenteral route is employed. Thus, for example, for intravenous administration, a dose in the range, for example, 0.5 mg to 30 mg per kg body weight will generally be used. Similarly, for administration by inhalation, a dose in the range, for example, 0.5 mg to 25 mg per kg body weight will be used. Oral administration is however preferred.

The following Examples illustrate the invention.

#### **Example 1 - Preparation of Compound No. 1 in Table 1**

A solution of 4-n-butoxyaniline (110 mg, 0.67 mmol) in isopropanol (7 ml) was added to 4-chloro-6,7-dimethoxyquinazoline hydrochloride (174 mg, 0.67 mmol) and the reaction heated at 73 °C for 2 hours before being cooled to 5 °C. The solid which precipitated was collected by suction filtration and washed with diethyl ether (2 x 5 ml). Drying of this material yielded the title compound (80 mg, 34 % yield) as an off-white solid :

<sup>1</sup>H-NMR (DMSO d<sub>6</sub>) : 11.25 (s, 1H), 8.74 (s, 1H), 8.24 (s, 1H), 7.54 (d, 2H, J = 8 Hz), 7.31 (s, 1H), 7.01 (d, 2H, J = 8 Hz), 3.99 (m, 2H), 3.99 (s, 3H), 3.97 (s, 3H), 1.70 (m, 2H), 1.45 (m, 2H), 0.93 (t, 3H, J = 7 Hz) :

MS (-ve ESI) : 352 (M-H)<sup>-</sup>,

MS (+ve ESI) : 354 (M+H)<sup>+</sup>.

4-Chloro-6,7-dimethoxyquinazoline, used as the starting material was obtained as follows :

a) A mixture of 4,5-dimethoxyanthranilic acid (19.7g, 100 mmol) and formamide (10ml) was heated at 190 °C for 5 hours. The mixture was allowed to cool to approximately 80 °C and water (50ml) was added. The mixture was then allowed to stand at ambient temperature

for 3 hours. Collection of the solid by suction filtration, followed by washing with water (2 x 50 ml) and drying *in vacuo*, yielded 6,7-dimethoxy-3,4-dihydroquinazolin-4-one (3.65g, 18 % yield) as a white solid.

<sup>1</sup>H-NMR (DMSO d<sub>6</sub>) : 12.10 (s, 1H), 7.95 (s, 1H), 7.42 (s, 1H), 7.11 (s, 1H), 3.88 (s, 3H), 3.84 (s, 3H) :

MS (-ve ESI) : 205 (M-H)<sup>-</sup>.

b) Dimethylformamide (0.2 ml) was added dropwise to a solution of 6,7-dimethoxy-3,4-dihydro-quinazolin-4-one (10.0 g, 48.5 mmol) in thionyl chloride (200ml) and the reaction was heated at reflux for 6 hours. The reaction was cooled, excess thionyl chloride was removed *in vacuo* and the residue was azeotroped with toluene (2 x 50 ml) to remove the last of the thionyl chloride. The residue was taken up in dichloromethane (550 ml), the solution was washed with saturated aqueous sodium hydrogen carbonate solution (2 x 250 ml) and the organic phase was dried over magnesium sulphate. Solvent evaporation *in vacuo* yielded 4-chloro-6,7-dimethoxyquinazoline (10.7 g, 98 % yield) as a white solid :

<sup>1</sup>H-NMR (DMSO d<sub>6</sub>) : 8.86 (s, 1H), 7.42 (s, 1H), 7.37 (s, 1H), 4.00 (s, 3H), 3.98 (s, 3H) :

MS (+ve ESI) : 225 (M-H)<sup>+</sup>.

### **Example 2 - Preparation of Compound No. 2 in Table 1**

2-Picolyl chloride hydrochloride (260 mg, 1.59 mmol) was added to a suspension of potassium carbonate (796 mg, 5.77 mmol), potassium iodide (358 mg, 2.16 mmol) and 4-(4-hydroxy-3-methylanilino)-6,7-dimethoxyquinazoline (500 mg, 1.92 mmol) in acetone (25 ml) and the reaction heated at reflux for 18 hours. The reaction was cooled, filtered and the filtrate evaporated *in vacuo*. Purification by flash chromatography on silica gel, eluting with 0-4% methanol in dichloromethane, yielded the title compound (436 mg, 68 % yield) as a white solid:

<sup>1</sup>H-NMR (DMSO d<sub>6</sub>) : 11.36 (s, 1H), 8.69 (d, 1H, J = 6 Hz), 8.30 (s, 1H), 8.09 (dt, 1H, J = 2,7 Hz), 7.74 (d, 2H, J = 8 Hz), 7.57 (m, 1H), 7.41-7.45 (m, 2H), 7.34 (s, 1H), 7.09 (d, 1H, J = 8 Hz), 5.34 (s, 2H), 3.99 (s, 3H), 3.97 (s, 3H), 2.28 (s, 3H) :

MS (-ve ESI) : 401 (M-H)<sup>-</sup>,

5 MS (+ve ESI) : 403 (M+H)<sup>+</sup>.

4-(4-hydroxy-3-methylanilino)-6,7-dimethoxyquinazoline, used as starting material, was obtained as follows:-

An analogous reaction to that described in example 1, but starting with 4-amino-2-methylphenol (6.98 g, 56.7 mmol) and 4-chloro-6,7-dimethoxyquinazoline hydrochloride  
10 (14.79 g, 56.7 mmol), yielded the title compound (17.72 g, 90 % yield) as a white solid :  
MS (+ve CI) : 312 (M+H)<sup>+</sup>.

### Example 3 - Preparation of Compound No. 3 in Table 1

An analogous reaction to that described in example 1, but starting with 3-methyl-4-((4-methyl-2-pyridyl)methoxy)aniline (400 mg, 1.5 mmol) yielded the title compound (294 mg, 47  
15 % yield) as an off-white solid :

<sup>1</sup>H-NMR (DMSO d<sub>6</sub>) : 11.36 (s, 1H), 8.76 (s, 1H), 8.62 (d, 1H, J = 7 Hz), 8.30 (s, 1H), 7.71 (s, 1H), 7.54 (d, 1H, J = 8 Hz), 7.42-7.46 (m, 2H), 7.34 (s, 1H), 7.10 (d, 1H, J = 8 Hz), 5.36 (s, 2H), 3.99 (s, 3H), 3.97 (s, 3H), 2.44 (s, 3H), 2.28 (s, 3H) :

20 MS (+ve ESI) : 417 (M+H)<sup>+</sup>.

3-Methyl-4-((4-methyl-2-pyridyl)methoxy)aniline, used as starting material, was obtained as follows :-

a) n-Butyllithium (24 ml of a 1.6 N solution in hexanes, 38.4 mmol) was added to a stirred solution of 2,4-lutidine (4.28 g, 40 mmol) in tetrahydrofuran (70 ml) at -70 °C under an  
25 inert atmosphere. After 1 hour, air was bubbled through (for 1 hour), methanol (50 ml) was added and the reaction allowed to warm to ambient temperature. The reaction mixture was filtered and then evaporated *in vacuo*. Purification of the crude product by flash chromatography on silica gel, eluting with ethyl acetate, yielded 2-(hydroxymethyl)-4-picoline (700 mg, 14 % yield) as a white solid.

30 b) Sodium hydride (150 mg of an 80% dispersion in mineral oil, 5.00 mmol) was added to a stirred solution of 2-(hydroxymethyl)-4-picoline (590 mg, 5.00 mmol) in N-

methylpyrrolidine (20ml) at ambient temperature. 2-Fluoro-5-nitrotoluene (775 mg, 5.00 mmol) was added, the reaction was stirred for 18 hours at ambient temperature and the reaction was poured into water (60 ml). Collection of the yellow solid which precipitated, followed by drying *in vacuo*, yielded 2-((4-methyl-2-pyridyl)methoxy)-5-nitrotoluene (900 mg, 70 % yield) as a yellow solid.

c) 5% Platinum on carbon (50 mg) was added to a solution of 2-((4-methyl-2-pyridyl)methoxy)-5-nitrotoluene (750 mg, 2.91 mmol) in ethanol (150 ml) and the solution was stirred for 2 hours at ambient temperature under an atmosphere of hydrogen. Filtration of the reaction mixture, followed by solvent evaporation *in vacuo*, yielded 3-methyl-4-((4-methyl-2-pyridyl)methoxy)aniline (420 mg, 63 % yield) as a yellow gum.

#### Example 4 - Preparation of Compound No. 4 in Table 1

An analogous reaction to that described in example 1, but starting with 3-methyl-4-((4-methoxy-2-pyridyl)methoxy)aniline (670 mg, 2.75 mmol) yielded the title compound (290 mg, 24 % yield) as a brown solid :

<sup>1</sup>H-NMR (DMSO d<sub>6</sub>) : 9.60 (s, 1H), 8.43 (s, 1H), 8.39 (d, 1H, J = 7 Hz), 7.85 (s, 1H), 7.47-7.49 (m, 2H), 7.15 (s, 1H), 7.02 (d, 1H, J = 2 Hz), 6.97 (d, 1H, J = 8 Hz), 6.92 (dd, 1H, J = 2,8 Hz), 5.14 (s, 2H), 3.93 (s, 3H), 3.91 (s, 3H), 3.83 (s, 3H), 2.27 (s, 3H) :

MS (-ve ESI) : 431 (M-H)<sup>-</sup>,

MS (+ve ESI) : 433 (M+H)<sup>+</sup>.

3-Methyl-4-((4-methoxy-2-pyridyl)methoxy)aniline, used as starting material, was obtained as follows :-

a) A solution of 2-picoline carboxylic acid (10.7 g, 87 mmol) in thionyl chloride (50 ml) was heated at reflux for 18 hours before being cooled and evaporated *in vacuo*. The residue was treated with methanol (25 ml) and then added to a solution of sodium methoxide prepared from sodium (1.0 g, 43 mmol) and methanol (100 ml). The reaction was heated at reflux for 3 hours, cooled and evaporated *in vacuo*. The residue was partitioned between water and ethyl acetate and the organic phase was separated. Evaporation of the organic phase yielded methyl 4-methoxypicoline-2-carboxylate (6.00 g, 41 % yield) as a white solid.

b) Lithium aluminum hydride (16 ml of a 1.0 N solution in diethyl ether, 16 mmol) was added to a solution of 2-(hydroxymethyl)-4-methoxypyridine (2.70 g, 16 mmol) in diethyl ether (50 ml) at ambient temperature. The reaction was stirred for 1 hour, poured into an aqueous solution of Rochelle's salt (250 ml) and the reaction mixture extracted with ethyl acetate (3 x 50 ml). Purification of the crude product by flash chromatography on silica gel, eluting with dichloromethane-ethyl acetate, yielded 2-(hydroxymethyl)-4-methoxypyridine (800 mg, 36 % yield) as a white solid.

c) An analogous reaction to that described in example 3b, but starting with 2-(hydroxymethyl)-4-methoxypyridine (600 mg, 4.30 mmol), yielded 2-((4-methoxy-2-pyridyl)methoxy)-5-nitrotoluene (780 mg, 70 % yield) as a yellow solid.

d) An analogous reaction to that described in example 3c, but starting with 2-((4-methoxy-2-pyridyl)methoxy)-5-nitrotoluene (770 mg, 2.96 mmol) yielded 3-methyl-4-((4-methoxy-2-pyridyl)methoxy)aniline (680 mg, 99 % yield) as a yellow solid.

#### Example 5 - Preparation of Compound No. 5 in Table 1

An analogous reaction to that described in example 1, but starting with 3-methyl-4-((6-methyl-2-pyridyl)methoxy)aniline (1.50 g, 6.14 mmol) yielded the title compound (748 mg, 29 % yield) as a white solid :

<sup>1</sup>H-NMR (DMSO d<sub>6</sub>) : 11.40 (s, 1H), 8.76 (s, 1H), 8.32 (s, 1H), 8.13 (t, 1H, J = 7 Hz), 7.67 (d, 1H, J = 8 Hz), 7.56 (d, 1H, J = 8 Hz), 7.43-7.47 (m, 2H), 7.35 (s, 1H), 7.09 (d, 1H, J = 8 Hz), 5.38 (s, 2H), 4.00 (s, 3H), 3.97 (s, 3H), 2.65 (s, 3H), 2.28 (s, 3H) :

MS (-ve ESI) : 415 (M-H)<sup>-</sup>,

MS (+ve ESI) : 417 (M+H)<sup>+</sup>.

3-Methyl-4-((4-methyl-2-pyridyl)methoxy)aniline, used as starting material, was obtained as follows :-

a) Sodium hydride (150 mg of an 80% dispersion in mineral oil, 5.00 mmol) was added to a stirred solution of 2-(hydroxymethyl)-4-picoline (590 mg, 5.00 mmol) in N-methylpyrrolidine (20ml) at ambient temperature. 2-Fluoro-5-nitrotoluene (775 mg, 5.00 mmol) was added, the reaction was stirred for 18 hours at ambient temperature and the reaction was poured into water (60 ml). Collection of the yellow solid which precipitated,

followed by drying *in vacuo*, yielded 2-((4-methyl-2-pyridyl)methoxy)-5-nitrotoluene (900 mg, 70 % yield) as a yellow solid.

a) An analogous reaction to that described in example 3b, but starting with 2-(hydroxymethyl)-6-methylpyridine (2.43 g, 20 mmol), yielded 2-((6-methyl-2-pyridyl)methoxy)-5-nitrotoluene (2.70 g, 52 % yield) as a yellow solid.

b) An analogous reaction to that described in example 3c, but starting with 2-((6-methyl-2-pyridyl)methoxy)-5-nitrotoluene (400 mg, 1.55 mmol) yielded 3-methyl-4-((6-methyl-2-pyridyl)methoxy)aniline (300 mg, 85 % yield) as a yellow gum.

#### **Example 6 - Preparation of Compound No. 6 in Table 1**

An analogous reaction to that described in example 1, but starting with 4-chloro-6-acetoxy-7-methoxyquinazoline (150 mg, 0.60 mmol) and 3-fluoro-4-(2-pyridylmethoxy)aniline (142 mg, 0.65 mmol) yielded the title compound (200 mg, 77 % yield) as a white solid :

<sup>1</sup>H-NMR (DMSO d<sub>6</sub>) : 11.06 (s, 1H), 8.85 (s, 1H), 8.59 (m, 2H), 7.88 (dt, 1H, J = 1,7 Hz), 7.72 (dd, 1H, J = 2,8 Hz), 7.56 (d, 1H, J = 8 Hz), 7.29-7.46 (m, 4H), 5.30 (s, 2H), 3.99 (s, 3H), 2.37 (s, 3H) :

MS (-ve ESI) : 433 (M-H)<sup>-</sup>,

MS (+ve ESI) : 435 (M+H)<sup>+</sup>.

4-chloro-6-acetoxy-7-methoxyquinazoline and 3-fluoro-4-(2-pyridylmethoxy)aniline, used as the starting materials, were obtained as follows :

a) A mixture of 6,7-dimethoxy-3,4-dihydroquinazolin-4-one (20.0 g, 97 mmol) and racemic methionine (21.7 g, 146 mmol) in methanesulphonic acid (150 ml) were heated at 100 °C for 5.5 hours and then allowed to cool to ambient temperature over 18 hours. The reaction was poured into cold water (750 ml), the pH of the aqueous solution was adjusted to pH 6 (by addition of 2.0N aqueous sodium hydroxide solution) and the solid which formed was collected by suction filtration. The solid was dried *in vacuo* and then dissolved in a mixture of pyridine (20 ml) and acetic anhydride (150 ml). The solution was heated at 100 °C for 1 hour, cooled and poured into cold water (1050 ml). Collection of the resultant solid by suction filtration, followed by drying *in vacuo*, yielded 6-acetoxy-7-methoxy-3,4-dihydroquinazolin-4-one (13.9 g, 57 % yield) as a pale-brown solid :



<sup>1</sup>H-NMR (DMSO d<sub>6</sub>) : 12.16 (s, 1H), 8.05 (s, 1H), 7.75 (s, 1H), 3.90 (s, 3H), 2.25 (s, 3H) :

MS (-ve ESI) : 233 (M-H)<sup>-</sup>,

b) Dimethylformamide (0.25 ml) was added dropwise to a solution of 6-acetoxy-7-methoxy-3,4-dihydro-quinazolin-4-one (13.8 g, 59.0 mmol) in thionyl chloride (150ml) and the reaction was heated at reflux for 1.5 hours. The reaction was cooled, excess thionyl

chloride was removed *in vacuo* and the residue was azeotroped with toluene (2 x 50 ml) to remove the last of the thionyl chloride. Drying *in vacuo* yielded 4-chloro-6,7-dimethoxyquinazoline hydrochloride (14.7 g, 87 % yield) as a beige solid, which was used without further purification :

<sup>1</sup>H-NMR (DMSO d<sub>6</sub>) : 9.0 (s, 1H), 8.0 (s, 1H), 7.6 (s, 1H), 4.0 (s, 3H), 2.35 (s, 3H) :

MS (+ve ESI) : 253 (M+H)<sup>+</sup>.

c) An analogous reaction to that described in example 3b, but starting with 2-(hydroxymethyl)pyridine (3.50 g, 36 mmol) and 3,4-difluoronitrobenzene (5.00 g, 31.4 mmol), yielded 2-((6-methyl-2-pyridyl)methoxy)-5-nitrofluorobenzene (4.50 g, 58 % yield) as a yellow solid.

d) An analogous reaction to that described in example 3c, but starting with 2-((6-methyl-2-pyridyl)methoxy)-5-nitrofluorobenzene (4.5 g, 18.1 mmol), yielded 3-fluoro-4-(2-pyridylmethoxy)aniline (1.86 g, 47 % yield) as a yellow solid.

#### **Example 7 - Preparation of Compound No. 7 in Table 1**

4-Chloro-6,7-dimethoxyquinazoline (112 mg, 0.50 mmol) and potassium carbonate (69 mg, 0.50 mmol) were added sequentially to a stirred suspension of 4-propoxyphenol (76 mg, 0.50 mmol) in dimethylformamide (3 ml). The reaction was heated at 100 °C for 4 hours then allowed to stir for a further 36 hours at ambient temperature. Brine (10 ml) was added and the reaction allowed to stand for 16 hours before the solid was collected by suction filtration (analogous reactions which failed to yield a solid precipitate were extracted with dichloromethane (2 x 5 ml) and the dichloromethane layer evaporated *in vacuo* to give a solid product). Drying *in vacuo* yielded the title compound (66.3 mg, 56 % yield) as a white solid :  
<sup>1</sup>H-NMR (DMSO d<sub>6</sub>) : 8.55 (s, 1H), 7.57 (s, 1H), 7.39 (s, 1H), 7.22 (d, 2H), 7.03 (d, 2H), 4.00 (s, 3H), 3.98 (s, 3H), 3.97 (t, 2H), 1.70-1.82 (m, 2H), 1.02 (t, 3H) :  
 MS (+ve ESI) : 341 (M+H)<sup>+</sup>.

**Example 8 - Preparation of Compound No. 8 in Table 1**

An analogous reaction to that described in example 7, but starting with 4-phenoxyphenol (85 mg, 0.50 mmol) yielded the title compound (165 mg, 84 % yield) as a white solid :

<sup>1</sup>H-NMR (DMSO d<sub>6</sub>) : 8.59 (s, 1H), 7.58 (s, 1H), 7.44 (t, 2H), 7.40 (s, 1H), 7.36 (d, 2H), 7.18 (t, 1H), 7.11 (d, 2H), 7.07 (d, 2H), 4.00 (s, 3H), 3.99 (s, 3H) :  
MS (+ve ESI) : 375 (M+H)<sup>+</sup>.

**Example 9 - Preparation of Compound No. 9 in Table 1**

An analogous reaction to that described in example 7, but starting with 4-benzyloxyphenol (100 mg, 0.50 mmol) yielded the title compound (182 mg, 94 % yield) as a white solid :

<sup>1</sup>H-NMR (400MHz, DMSO d<sub>6</sub>) : 8.54 (s, 1H), 7.57 (s, 1H), 7.50 (d, 2H), 7.43 (t, 2H), 7.40 (s, 1H), 7.35 (t, 1H), 7.25 (d, 2H), 7.11 (d, 2H), 5.16 (s, 2H), 3.99 (s, 3H), 3.98 (s, 3H) :  
MS (+ve ESI) : 387 (M+H)<sup>+</sup>.

**Example 10 - Preparation of Compound No. 10 in Table 1**

An analogous reaction to that described in example 7, but starting with 4-(methylmercapto)phenol (70 mg, 0.50 mmol) yielded the title compound (146 mg, 89 % yield) as a white solid :

<sup>1</sup>H-NMR (DMSO d<sub>6</sub>) : 8.55 (s, 1H), 7.56 (s, 1H), 7.36-7.42 (m, 3H), 7.29 (d, 2H), 3.99 (s, 3H), 3.98 (s, 3H), 2.53 (s, 3H) :  
MS (+ve ESI) : 329 (M+H)<sup>+</sup>.

**Example 11 - Preparation of Compound No. 11 in Table 1**

An analogous reaction to that described in example 7, but starting with 4-pentyloxyphenol (90 mg, 0.50 mmol) yielded the title compound (166 mg, 90 % yield) as a white solid :

<sup>1</sup>H-NMR (DMSO d<sub>6</sub>) : 8.54 (s, 1H), 7.56 (s, 1H), 7.38 (s, 1H), 7.21 (d, 2H), 7.02 (d, 2H), 4.01 (t, 2H), 4.00 (s, 3H), 3.99 (s, 3H), 1.71-1.81 (m, 2H), 1.32-1.49 (m, 4H), 0.92 (t, 3H) :

MS (+ve ESI) : 369 (M+H)<sup>+</sup>.

**Example 12 - Preparation of Compound No. 12 in Table 1**

A mixture of 4-(4-hydroxyanilino)-6,7-dimethoxyquinazoline hydrochloride (100 mg, 0.30 mmol), potassium carbonate (137 mg, 0.99 mmol) and 2-picolyl chloride hydrochloride (54 mg, 0.33 mmol) were heated in dimethylformamide (5 ml) at 100 °C for 4 hours and then allowed to cool to ambient temperature. The reaction was poured into water (50 ml) and the aqueous phase was extracted with dichloromethane (3 x 50 ml). The combined organic layers were evaporated *in vacuo* to a volume of 10 ml, and then diethyl ether was added (25 ml) to cause precipitation of a brown solid. Purification by chromatography on a silica gel bond-elute cartridge, eluting with 4% methanol in dichloromethane yielded the title compound (48 mg, 41 % yield) as a white solid :

<sup>1</sup>H-NMR (DMSO d<sub>6</sub>) : 9.36 (s, 1H), 8.58 (d, 1H, J = 8 Hz), 8.36 (s, 1H), 7.79-7.85 (m, 1H), 7.79 (s, 1H), 7.61 (d, 2H, J = 8 Hz), 7.52 (d, 1H, J = 8 Hz), 7.31-7.35 (m, 1H), 7.14 (s, 1H), 7.04 (d, 2H, J = 8 Hz), 5.19 (s, 2H), 3.94 (s, 3H), 3.88 (s, 3H) :

MS (-ve ESI) : 236 (M-H)<sup>-</sup>,

MS (+ve ESI) : 238 (M+H)<sup>+</sup>.

4-(4-Hydroxyanilino)-6,7-dimethoxyquinazoline hydrochloride, used as the starting material was obtained as follows :

A mixture of 4-chloro-6,7-dimethoxyquinazoline (1.00 g, 4.45 mmol and 4-aminophenol (530 mg, 4.90 mmol) were heated in isopropanol (50 ml) at reflux for 2 hours and then allowed to cool to ambient temperature. Collection of the precipitated solid by suction filtration, followed by washing with diethyl ether and drying *in vacuo*, yielded 4-(4-hydroxyanilino)-6,7-dimethoxyquinazoline hydrochloride (1.34 g, 90 % yield) as a white solid :

<sup>1</sup>H-NMR (DMSO d<sub>6</sub>) : 11.24 (s, 1H), 9.66 (s, 1H), 8.72 (s, 1H), 8.24 (s, 1H), 7.40 (d, 2H, J = 8 Hz), 7.34 (s, 1H), 6.84 (d, 2H, J = 8 Hz), 3.96 (s, 3H), 3.94 (s, 3H) :

MS (-ve ESI) : 296 (M-H)<sup>-</sup>,

MS (+ve ESI) : 298 (M+H)<sup>+</sup>.

**Example 13 - Preparation of Compound No. 13 in Table 1**

An analogous reaction to that described in example 12, but starting with phenethyl bromide (90.8 mg, 0.40 mmol), potassium carbonate (96 mg, 0.69 mmol) and 4-(4-hydroxyanilino)-6,7-dimethoxyquinazoline hydrochloride (105 mg, 0.31 mmol), yielded the title compound (41 mg, 33 % yield) as a pale yellow solid, after purification by chromatography on silica gel, eluting with 2% methanol in dichloromethane :

<sup>1</sup>H-NMR (DMSO d<sub>6</sub>) : 8.61 (s, 1H), 7.51 (d, 2H, J = 8 Hz), 7.23-7.35 (m, 6H), 7.05 (s, 1H), 6.99 (s, 1H), 6.94 (d, 2H, J = 8 Hz), 4.20 (t, 2H, J = 8 Hz), 4.01 (s, 3H), 3.98 (s, 3H), 3.11 (t, 2H, J = 8 Hz) :

MS (-ve ESI) : 400 (M-H)<sup>-</sup>,

MS (+ve ESI) : 402 (M+H)<sup>+</sup>.

**Example 14 - Preparation of Compound No. 14 in Table 1**

An analogous reaction to that described in example 12, but starting with allyl bromide (0.055 ml, 0.64 mmol), potassium carbonate (96 mg, 0.69 mmol) and 4-(4-hydroxyanilino)-6,7-dimethoxyquinazoline hydrochloride (105 mg, 0.31 mmol), yielded the title compound (42 mg, 39 % yield) as a pale yellow solid, after purification by chromatography on silica gel, eluting with 2% methanol in dichloromethane :

<sup>1</sup>H-NMR (DMSO d<sub>6</sub>) : 8.61 (s, 1H), 7.51 (d, 2H, J = 8 Hz), 7.24 (s, 1H), 7.12 (s, 1H), 7.01 (s, 1H), 6.96 (d, 2H, J = 8 Hz), 6.00-6.14 (m, 1H), 5.43 (dd, 1H, J = 2, 16 Hz), 5.28 (dd, 1H, J = 2, 10 Hz), 4.54 (d, 2H, J = 7 Hz), 4.00 (s, 3H), 3.96 (s, 3H) :

MS (-ve ESI) : 336 (M-H)<sup>-</sup>,

MS (+ve ESI) : 338 (M+H)<sup>+</sup>.

**Example 15 - Preparation of Compound No. 15 in Table 1**

An analogous reaction to that described in example 1, but starting with 4-chloro-6-methoxy-7-(3-morpholinopropoxy)quinazoline (74 mg, 0.22 mmol) and (4-amino-2,6-dichlorophenyl)-4-chlorophenylether (70 mg, 0.24 mmol) yielded the title compound (115 mg, 86 % yield) as a white solid :

<sup>1</sup>H-NMR (DMSO d<sub>6</sub>) : 8.82 (s, 1H), 8.35 (s, 1H), 8.11 (d, 1H), 7.79 (dd, 1H), 7.43 (d, 2H), 7.38 (s, 1H), 7.28 (d, 1H), 7.00 (d, 2H), 4.32 (t, 2H), 4.02 (s, 3H), 3.99 (m, 2H), 3.80 (m, 2H), 3.48 (m, 2H), 3.30 (m, 2H), 3.11 (m, 2H), 2.30 (m, 2H) ;  
MS (+ve ESI) : 555 (M+H)<sup>+</sup>.

- 5 4-Chloro-6-methoxy-7-(3-morpholinopropoxy)quinazoline, used as the starting material, was obtained as follows :

a) A mixture of morpholine (261 ml, 3.00 mol) and 1-bromo-3-chloropropane (148 ml, 1.50 mol) in toluene (900 ml) was stirred for 18 hours at ambient temperature. Additional 1-bromo-3-chloropropane (25 ml, 0.25 mol) was added, the reaction was stirred for a further 1  
10 hour and then filtered to remove the precipitated solid before the filtrate was concentrated *in vacuo*. Distillation of the crude oil yielded N-(3-chloropropyl)-morpholine (119.3 g, 49 % yield) as the fraction boiling at 70 - 80 °C / 2.6 mmHg :

<sup>1</sup>H-NMR (DMSO d<sub>6</sub>) : 3.65 (t, 2H), 3.55 (m, 4H), 2.4 (t, 2H), 2.39 (m, 4H), 1.85 (m, 2H) :  
MS (+ve ESI) : 164 (M+H)<sup>+</sup>.

15 b) N-(3-Chloropropyl)morpholine (90 g, 0.55 mol) was added dropwise, over 30 minutes, to a solution of ethyl vanillate (98 g, 0.50 mol) and powdered potassium carbonate (104 g, 0.75 mol) in dimethylformamide (300 ml) at 80 °C. The reaction was heated at 80 °C for 90 minutes, cooled to ambient temperature, filtered and the filtrate concentrated *in vacuo*. The crude product was taken up in diethyl ether (1000 ml), filtered and washed with water (2 x 200  
20 ml) and brine (200 ml). Solvent evaporation in *vacuo* yielded ethyl 3-methoxy-4-(3-morpholinopropoxy)benzoate (161.5 g, 100 % yield) as a pale yellow oil which crystallised on standing to afford a pale yellow solid :

<sup>1</sup>H-NMR (DMSO d<sub>6</sub>) : 7.55 (dd, 1H), 7.4 (d, 1H), 7.05 (d, 1H), 4.3 (q, 2H), 4.05 (t, 2H), 3.8 (s, 3H), 3.55 (m, 4H), 2.4 (t, 2H), 2.35 (m, 4H), 1.9 (m, 2H), 1.3 (t, 3H) :

25 MS (-ve ESI) : 324 (M-H)<sup>-</sup>,

c) Concentrated sulphuric acid (110 ml) and concentrated nitric acid (19.0 ml, 0.289 mol) were added cautiously, over a 50 minute period, to a two-phase system containing a stirred solution of ethyl 3-methoxy-4-(3-morpholinopropoxy)benzoate (76.5 g, 0.237 mol) in dichloromethane (600 ml), acetic acid (300 ml) and water (70 ml) at 5 °C. The reaction was  
30 allowed to warm to ambient temperature over 18 hours, the aqueous phase was separated, and the aqueous phase was taken to pH 9 by addition of 40% aqueous sodium hydroxide solution

(775 ml). Extraction of the aqueous phase with dichloromethane (3 x 600 ml) and subsequent solvent evaporation *in vacuo* yielded ethyl 3-methoxy-4-(3-morpholinopropoxy)-6-nitrobenzoate (141.3 g, 86 % yield) as a yellow gum :

<sup>1</sup>H-NMR (CDCl<sub>3</sub>) : 7.5 (s, 1H), 7.1 (s, 1H), 4.4 (q, 2H), 4.2 (t, 2H), 4.0 (s, 3H), 3.7 (m, 4H),  
5 2.5 (t, 2H), 2.45 (m, 4H), 2.05 (m, 2H), 1.4 (t, 3H) :

MS (+ve ESI) : 369 (M+H)<sup>+</sup>.

d) A suspension of ethyl 3-methoxy-4-(3-morpholinopropoxy)-6-nitrobenzoate (132.2 g, 359 mmol) and 10% palladium on carbon (3.0 g) in a mixture of ethanol (200 ml) and ethyl acetate (2000 ml) was stirred under an atmosphere of hydrogen for 18 hours. Removal of the  
10 catalyst by filtration, followed by solvent evaporation *in vacuo* yielded ethyl 3-methoxy-4-(3-morpholinopropoxy)-6-aminobenzoate (122 g, 100 % yield) as a brown oil :

<sup>1</sup>H-NMR (DMSO d<sub>6</sub>) : 7.15 (s, 1H), 6.4 (s, 2H), 6.35 (s, 1H), 4.2 (q, 2H), 3.95 (t, 2H), 3.65 (s, 3H), 3.55 (m, 4H), 2.4 (t, 2H), 2.35 (m, 4H), 1.85 (m, 2H), 1.25 (t, 3H) :

MS (-ve ESI) : 337 (M-H)<sup>-</sup>,

15 MS (+ve ESI) : 339 (M+H)<sup>+</sup>.

e) A solution of ethyl 3-methoxy-4-(3-morpholinopropoxy)-6-aminobenzoate (130 g, 384 mmol) in formamide (280 ml) was heated at 180 °C for 3 hours, during which time a small amount (25 ml) of liquid distilled out of the reaction. The reaction was cooled to 125 °C and the excess formamide was evaporated *in vacuo*. Trituration of the solid residue with  
20 isopropanol (100 ml), followed by drying *in vacuo*, yielded 6-methoxy-7-(3-morpholinopropoxy)-3,4-dihydroquinazolin-4-one (83.0 g, 68 % yield) as a pale brown solid :

<sup>1</sup>H-NMR (DMSO d<sub>6</sub>) : 12.0 (s, 1H), 7.95 (s, 1H), 7.45 (s, 1H), 7.1 (s, 1H), 4.15 (t, 2H), 3.85 (s, 3H), 3.6 (m, 4H), 2.45 (t, 2H), 2.35 (m, 4H), 1.9 (m, 2H) :

MS (-ve ESI) : 318 (M-H)<sup>-</sup>,

25 MS (+ve ESI) : 320 (M+H)<sup>+</sup>.

f) Dimethylformamide (2.0 ml) was added dropwise to a solution of 6-methoxy-7-(3-morpholinopropoxy)-3,4-dihydroquinazolin-4-one (83.0 g, 261 mmol) in thionyl chloride (700ml) and the reaction was heated at reflux for 3.5 hours. The reaction was cooled, excess thionyl chloride was removed *in vacuo*, the residue was taken up in water (500 ml) and this  
30 aqueous solution was taken to pH 9 by addition of saturated aqueous sodium bicarbonate solution (300 ml). The aqueous phase was extracted with dichloromethane (2 x 400 ml), the

organic solution was washed with brine (400 ml) and the solvents were removed *in vacuo*. Trituration of the solid residue with ethyl acetate (150 ml), followed by drying *in vacuo*, yielded 4-chloro-6-methoxy-7-(3-morpholinopropoxy)quinazoline (53 g, 60 % yield) as a pale brown solid :

5 <sup>1</sup>H-NMR (CDCl<sub>3</sub>) : 8.85 (s, 1H), 7.39 (s, 1H), 7.38 (s, 1H), 4.3 (t, 2H), 4.05 (s, 3H), 3.7 (m, 4H), 2.6 (t, 2H), 2.5 (m, 4H), 2.1 (m, 2H) :  
MS (+ve ESI) : 338 (M+H)<sup>+</sup>.

#### **Example 16 - Preparation of Compound No. 16 in Table 1**

10 An analogous reaction to that described in example 15, but starting with (4-amino-2,6-dichlorophenyl)-4-chlorophenylsulphide (73 mg, 0.24 mmol) yielded the title compound (118 mg, 86 % yield) as a white solid :

<sup>1</sup>H-NMR (DMSO d<sub>6</sub>) : 8.92 (s, 1H), 8.41 (s, 1H), 8.38 (s, 2H), 7.40 (s, 1H), 7.39 (d, 2H), 7.10 (d, 2H), 4.30 (t, 2H), 4.03 (s, 3H), 4.00 (m, 2H), 3.80 (m, 2H), 3.50 (m, 2H), 3.28 (m, 2H),  
15 3.10 (m, 2H), 2.30 (m, 2H) ;  
MS (-ve ESI) : 603 (M-H)<sup>-</sup>.

#### **Example 17 - Preparation of Compound No. 17 in Table 1**

20 An analogous reaction to that described in example 15, but starting 4-aminothioanisole (33 mg, 0.24 mmol), yielded the title compound (103 mg, 95 % yield) as a white solid :

<sup>1</sup>H-NMR (DMSO d<sub>6</sub>) : 8.77 (s, 1H), 8.30 (s, 1H), 7.65 (d, 2H), 7.32-7.40 (m, 3H), 4.30 (t, 2H), 4.01 (s, 3H), 3.72-4.01 (m, 4H), 3.00-3.54 (m, 6H), 2.54 (s, 3H), 2.22-2.38 (m, 2H) ;  
MS (+ve ESI) : 441 (M+H)<sup>+</sup>.

#### **Example 18 - Preparation of Compound No. 18 in Table 1**

25 An analogous reaction to that described in example 1, but starting with 4-benzyloxyaniline hydrochloride (118 mg, 0.50 mmol) and 4-chloro-6-methoxy-7-(3-morpholinopropoxy)quinazoline (168 g, 0.50 mmol), yielded the title compound (216 mg, 86 % yield) as a white solid :

<sup>1</sup>H-NMR (DMSO d<sub>6</sub>) : 9.8 (s, 1H), 8.3 (s, 1H), 7.7 (d, 2H), 7.4 (t, 2H), 7.35 (s, 1H), 7.15 (t, 1H), 7.1 (d, 2H), 7.05 (d, 2H), 4.3 (t, 2H), 4.0 (s, 3H), 3.95 (m, 2H), 3.8 (m, 2H), 3.5 (m, 2H), 3.3 (m, 2H), 3.1 (m, 2H), 2.3 (m, 2H) :

MS (+ve ESI) : 464 (M+H)<sup>+</sup>.

5

---

**Example 19 - Preparation of Compound No. 19 in Table 1**

A solution of 4-chloro-6-methoxy-7-benzyloxyquinazoline (150 mg, 0.50 mmol) and 4-phenoxyaniline (93 mg, 0.50 mmol), in isopropanol (5.0 ml) was at 40 °C for 30 minutes and then at 83 °C for 12 hours before the reaction was allowed to cool to ambient temperature.

10 The solid which had precipitated was collected by suction filtration and washed with diethyl ether (2 x 10 ml). Drying of this material yielded the title compound (209 mg, 86 % yield) as an off-white solid :

<sup>1</sup>H-NMR (DMSO d<sub>6</sub>) : 11.20 (s, 1H), 8.77 (s, 1H), 8.23 (s, 1H), 7.67 (d, 2H), 7.50 (d, 2H), 7.40-7.45 (m, 6H), 7.15 (d, 1H), 7.01-7.10 (m, 4H), 5.34 (s, 2H), 4.0 (s, 3H) :

15 MS (+ve ESI) : 450 (M+H)<sup>+</sup>.

4-Chloro-6-methoxy-7-benzyloxyquinazoline, used as the starting material, was obtained as follows :

a) A mixture of 2-amino-4-benzyloxy-5-methoxybenzamide (10g, 0.04mol), (prepared according to *J. Med. Chem.* 1977, 20, 146-149), and Gold's reagent (7.4g, 0.05mol) in dioxane  
20 (100ml) was stirred and heated at reflux for 24 hours. Sodium acetate (3.02g, 0.037mol) and acetic acid (1.65ml, 0.029mol) were added to the reaction mixture and it was heated for a further 3 hours. The volatiles were removed by evaporation, water was added to the residue, the solid was collected by filtration, washed with water and dried. Recrystallisation from acetic acid yielded 7-benzyloxy-6-methoxy-3,4-dihydroquinazolin-4-one (8.7g, 84 % yield) as  
25 a white solid:

b) Dimethylformamide (0.2 ml) was added dropwise to a solution of 6-methoxy-7-benzyloxy-3,4-dihydroquinazolin-4-one (5.00 g, 17.9 mmol) in thionyl chloride (100ml) and the reaction was heated at reflux for 1 hour. The reaction was cooled, excess thionyl chloride was removed *in vacuo* and the residue was azeotroped with toluene (3 x 50 ml) to remove the  
30 last of the thionyl chloride. The residue was taken up in dichloromethane (550 ml), the solution was washed with saturated aqueous sodium hydrogen carbonate solution (100 ml) and



water (100 ml) and the organic phase was dried over magnesium sulphate. Solvent evaporation *in vacuo* yielded 4-chloro-6,7-dimethoxyquinazoline (4.80 g, 90 % yield) as a pale brown solid :

<sup>1</sup>H-NMR (DMSO d<sub>6</sub>) : 8.85 (s, 1H), 7.58 (s, 1H), 7.5 (d, 2H), 7.4 (m, 4H), 5.35 (s, 2H), 4.0 (s, 3H) :

MS (+ve ESI) : 301 (M+H)<sup>+</sup>.

#### **Example 20 - Preparation of Compound No. 20 in Table 1**

A solution of 1.0N hydrochloric acid in ether (0.50 ml, 0.50 mmol) was added to a solution of 4-benzyloxyaniline hydrochloride (118 mg, 0.50 mmol) and 4-chloro-6-methoxy-7-(3-morpholinopropoxy)quinazoline (168 mg, 0.50 mmol), in isopropanol (5.0 ml). The reaction was heated at 40 °C for 30 minutes and then at 83 °C for 12 hours. The reaction was allowed to cool to ambient temperature and the solid which had precipitated was collected by suction filtration and washed with diethyl ether (2 x 10 ml). Drying of this material yielded the title compound (228 mg, 85 % yield) as a white solid :

<sup>1</sup>H-NMR (DMSO d<sub>6</sub>) : 15.00 (s, 1H), 11.34 (s, 1H), 11.12 (s, 1H), 8.75 (s, 1H), 8.33 (s, 1H), 7.59 (d, 2H), 7.30-7.52 (m, 6H), 7.12 (d, 2H), 5.16 (s, 1H), 4.30 (t, 2H), 4.01 (s, 3H), 3.73-4.01 (m, 4H), 2.92-3.58 (m, 6H), 2.21-2.39 (m, 2H) ;

MS (+ve ESI) : 501 (M+H)<sup>+</sup>.

#### **Example 21 - Preparation of Compound No. 21 in Table 1**

An analogous reaction to that described in example 20, but starting with 4-amino-4'-nitrodiphenylsulphide (123 mg, 0.50 mmol) yielded the title compound (281 mg, 96 % yield) as a white solid :

<sup>1</sup>H-NMR (DMSO d<sub>6</sub>) : 11.50 (s, 1H), 11.10 (s, 1H), 8.85 (s, 1H), 8.48 (s, 1H), 8.17 (d, 2H), 8.00 (d, 2H), 7.70 (d, 2H), 7.50 (s, 1H), 7.35 (d, 2H), 4.32 (t, 2H), 4.05 (s, 3H), 3.99 (m, 2H), 3.85 (m, 2H), 3.50 (m, 2H), 3.30 (m, 2H), 3.10 (m, 2H), 2.32 (m, 2H) ;

MS (+ve ESI) : 548 (M+H)<sup>+</sup>.

**Example 22 - Preparation of Compound No. 22 in Table 1**

An analogous reaction to that described in example 20, but starting with 4-butoxyaniline (82 mg, 0.50 mmol) yielded the title compound (237 mg, 94 % yield) as a white solid :

5 <sup>1</sup>H-NMR (DMSO d<sub>6</sub>) : 11.35 (s, 1H), 11.12 (s, 1H), 8.75 (s, 1H), 8.35 (s, 1H), 7.60 (d, 2H), 7.40 (s, 1H), 7.05 (d, 2H), 4.31 (t, 2H), 4.03 (m, 2H), 4.02 (s, 3H), 3.99 (m, 2H), 3.85 (m, 2H), 3.50 (m, 2H), 3.30 (m, 2H), 3.10 (m, 2H), 2.35 (m, 2H), 1.70 (m, 2H), 1.58 (m, 2H), 0.95 (t, 3H) ;

MS (+ve ESI) : 467 (M+H)<sup>+</sup>.

**Example 23 - Preparation of Compound No. 23 in Table 1**

An analogous reaction to that described in example 20, but starting with 4-amino-4'-chlorodiphenyl ether (110 mg, 0.50 mmol) yielded the title compound (244 mg, 88 % yield) as a white solid :

15 <sup>1</sup>H-NMR (DMSO d<sub>6</sub>) : 11.50 (s, 1H), 11.10 (s, 1H), 8.80 (s, 1H), 8.40 (s, 1H), 7.75 (d, 2H), 7.47 (d, 2H), 7.40 (s, 1H), 7.15 (d, 2H), 7.08 (d, 2H), 4.35 (t, 2H), 4.03 (s, 3H), 3.95 (m, 2H), 3.85 (m, 2H), 3.50 (m, 2H), 3.30 (m, 2H), 3.10 (m, 2H), 2.35 (m, 2H) ;

MS (+ve ESI) : 521 (M+H)<sup>+</sup>.

**Example 24 - Preparation of Compound No. 24 in Table 1**

20 An analogous reaction to that described in example 1, but starting with 4-chloro-6-methoxy-7-(2,2,2-trifluoroethoxy)quinazoline (572 mg, 1.96 mmol) and 3-fluoro-4-(2-pyridylmethoxy)-aniline (469 mg, 2.15 mmol) yielded the title compound (315 mg, 34 % yield) as a white solid :

25 <sup>1</sup>H-NMR (DMSO d<sub>6</sub>) : 11.59 (s, 1H), 8.83 (s, 1H), 8.65 (d, 2H, J = 5 Hz), 7.88 (dt, 1H, J = 1,7 Hz), 7.72 (dd, 1H, J = 2,8 Hz), 7.56 (d, 1H, J = 8 Hz), 7.29-7.46 (m, 4H), 5.30 (s, 2H), 5.05 (q, 2H, J = 8 Hz), 4.04 (s, 3H) :

MS (-ve ESI) : 473 (M-H)<sup>-</sup>,

MS (+ve ESI) : 475 (M+H)<sup>+</sup>.

30 4-Chloro-6-methoxy-7-(2,2,2-trifluoroethoxy)quinazoline, used as starting material was obtained as follows :

a) Potassium carbonate (62.2 g, 450 mmol) was added to a solution of ethyl vanillate (58.9 g, 300 mmol) in dimethylformamide (400 ml) and the reaction heated to 120 °C. 2,2,2-Trifluoroethyl methanesulphonate (63.4 g, 360 mmol) was added over 15 minutes and the reaction heated at 120 °C for 15 hours. The reaction was cooled to ambient temperature,  
5 diethyl ether (400 ml) was added and the reaction was filtered. The filtrate was evaporated *in*

*vacuo* and the residue was taken up in a mixture of diethyl ether (375 ml) and isohexane (375 ml). The organic layer was concentrated in *vacuo* to a total volume of 250 ml and the solid which crystallised out was collected by suction filtration. Drying of the solid in *vacuo* yielded ethyl 4-(2,2,2-trifluoroethoxy)-3-methoxybenzoate (43.0 g, 52 % yield) as a white crystalline  
10 solid :

<sup>1</sup>H-NMR (DMSO *d*<sub>6</sub>) : 7.57 (dd, 1H, J = 2, 8 Hz), 7.49 (d, 1H, J = 2 Hz), 7.18 (d, 1H, J = 8 Hz), 5.81 (q, 2H, J = 7 Hz), 5.29 (q, 2H, J = 7 Hz), 3.82 (s, 3H), 1.30 (t, 3H, J = 7 Hz) :  
MS (+ve ESI) : 279 (M+H)<sup>+</sup>.

b) Concentrated sulphuric acid (64 ml) and concentrated nitric acid (10.0 ml, 0.152 mol) were added cautiously, over 1 hour, to a two-phase system containing a stirred solution yielded  
15 ethyl 4-(2,2,2-trifluoroethoxy)-3-methoxybenzoate (35.3 g, 0.127 mol) in dichloromethane (340 ml), acetic acid (173 ml) and water (40 ml) at 5 °C. The reaction was allowed to warm to ambient temperature over 60 hours (with vigorous mechanical stirring), the aqueous phase was separated, and the organic phase washed with water (6 x 250 ml). The organic phase was  
20 concentrated to a total volume of ~200 ml, isohexane (150 ml) was added and the solid which precipitated out was collected by suction filtration. Drying of the solid *in vacuo* yielded ethyl 3-methoxy-4-(2,2,2-trifluoroethoxy)-6-nitrobenzoate (21.7 g, 52 % yield) as a yellow solid. The mother liquors contained a mixture of product (28%) and starting material (72%) which was recycled in a latter reaction :

25 <sup>1</sup>H-NMR (DMSO *d*<sub>6</sub>) : 7.80 (s, 1H), 7.42 (s, 1H), 4.90 (q, 2H, J = 7 Hz), 4.20-4.35 (m, 2H), 4.00 (s, 3H), 1.32 (t, 3H, J = 7 Hz) :  
MS (+ve ESI) : 324 (M+H)<sup>+</sup>.

c) A suspension of ethyl 3-methoxy-4-(2,2,2-trifluoroethoxy)-6-nitrobenzoate (24.0 g, 74.3 mmol) and 10% palladium on carbon (3.0 g) in a mixture of ethanol (100 ml) and ethyl  
30 acetate (750 ml) was stirred under an atmosphere of hydrogen for 18 hours. Removal of the

catalyst by filtration, followed by solvent evaporation *in vacuo* yielded ethyl 3-methoxy-4-(2,2,2-trifluoroethoxy)-6-aminobenzoate (20.2 g, 93 % yield) as a pale brown solid :

<sup>1</sup>H-NMR (DMSO *d*<sub>6</sub>) : 7.20 (s, 1H), 6.45 (s, 1H), 6.40 (s, 2H), 5.70 (q, 2H, J = 7 Hz), 4.20 (q, 2H, J = 7 Hz), 3.65 (s, 3H), 1.32 (t, 3H, J = 7 Hz) :

5 MS (-ve ESI) : 292 (M-H)<sup>-</sup>,

MS (+ve ESI) : 294 (M+H)<sup>+</sup>.

d) A mixture of ethyl 2-amino-4-(2,2,2-trifluoroethoxy)-5-methoxybenzoate (20.2 g, 69.1 mmol) and formamide (50ml) was heated at 175 °C for 6 hours. The mixture was allowed to cool to ambient temperature, ethanol (150 ml) was added and the reaction allowed to stand for 18 hours. Collection of the solid which had precipitated by suction filtration, followed by washing with ethanol (2 x 50 ml) and drying *in vacuo*, yielded 6-methoxy-7-(2,2,2-trifluoroethoxy)-3,4-dihydroquinazolin-4-one (15.8 g, 84 % yield) as a pale brown crystalline solid :

15 <sup>1</sup>H-NMR (DMSO *d*<sub>6</sub>) : 12.10 (s, 1H), 8.00 (s, 1H), 7.51 (s, 1H), 7.30 (s, 1H), 4.90 (q, 2H, J = 7 Hz), 3.90 (s, 3H) :

MS (-ve ESI) : 273 (M-H)<sup>-</sup>,

MS (+ve ESI) : 275 (M+H)<sup>+</sup>.

e) Dimethylformamide (0.1 ml) was added dropwise to a solution yielded 6-methoxy-7-(2,2,2-trifluoroethoxy)-3,4-dihydroquinazolin-4-one (15.8 g, 57.7 mmol) in thionyl chloride (200ml) and the reaction was heated at reflux for 6 hours. The reaction was cooled, excess thionyl chloride was removed *in vacuo* and the residue was azeotroped with toluene (2 x 50 ml) to remove the last of the thionyl chloride. The residue was taken up in dichloromethane (550 ml), the solution was washed with saturated aqueous sodium hydrogen carbonate solution (2 x 250 ml) and the organic phase was dried over magnesium sulphate. Solvent evaporation *in vacuo* yielded 4-chloro-6-methoxy-7-(2,2,2-trifluoroethoxy)quinazoline (16.3 g, 97 % yield) as a cream solid :

25 <sup>1</sup>H-NMR (DMSO *d*<sub>6</sub>) : 8.95 (s, 1H), 7.65 (s, 1H), 7.25 (s, 1H), 5.05 (q, 2H, J = 7 Hz), 4.00 (s, 3H) :

MS (+ve ESI) : 293, 295 (M+H)<sup>+</sup>.

**Example 25 - Preparation of Compound No. 25 in Table 1**

Triethylamine (0.10 ml, 0.72 mmol), tributylphosphine (0.45 ml, 1.83 mmol) and furfuryl alcohol (0.106 ml, 1.22 mmol) were added to a suspension of 4-(4-hydroxyanilino)-  
5 6,7-dimethoxyquinazoline hydrochloride (205 mg, 0.61 mmol) in dichloromethane (20 ml) at ambient temperature. The reaction was stirred for 20 minutes before addition of 1,1'-

(azodicarbonyl)dipiperidine (462 mg, 1.83 mmol) and then stirred for a further 3 hours.

Tributylphosphine (0.45 ml, 1.83 mmol) and 1,1'-(azodicarbonyl)dipiperidine (462 mg, 1.83 mmol) were added and the reaction stirred for 2 hours at ambient temperature. The reaction  
10 mixture was transferred to an SCX column which was eluted with 0-5% methanol in dichloromethane before the product was eluted with 3% ammonium hydroxide / 20% methanol in dichloromethane. Evaporation of the desired fractions *in vacuo*, followed by trituration of the solid product with ethyl acetate, yielded the title compound (34 mg, 15 % yield) as a white solid, after drying *in vacuo* :

15 <sup>1</sup>H-NMR (DMSO d<sub>6</sub>) : 9.38 (s, 1H), 8.39 (s, 1H), 7.81 (s, 1H), 7.69 (s, 1H), 7.63 (d, 1H, J = 8 Hz), 7.16 (s, 1H), 7.05 (d, 2H, J = 8 Hz), 6.58 (d, 1H, J = 5 Hz), 6.46 (d, 2H, J = 5 Hz), 5.07 (s, 2H), 3.96 (s, 3H), 3.93 (s, 3H) :

MS (-ve ESI) : 376 (M-H)<sup>-</sup>,

MS (+ve ESI) : 378 (M+H)<sup>+</sup>.

**Biological Data**

The compounds of the invention inhibit the serine/threonine kinase activity of the aurora2 kinase and thus inhibit the cell cycle and cell proliferation. These properties may be assessed, for example, using one or more of the procedures set out below:

**(a) In Vitro aurora2 kinase inhibition test**

This assay determines the ability of a test compound to inhibit serine/threonine kinase activity. DNA encoding aurora2 may be obtained by total gene synthesis or by cloning. This DNA may then be expressed in a suitable expression system to obtain polypeptide with serine/threonine kinase activity. In the case of aurora2, the coding sequence was isolated from  
30 cDNA by polymerase chain reaction (PCR) and cloned into the BamH1 and Not1 restriction endonuclease sites of the baculovirus expression vector pFastBac HTc (GibcoBRL/Life technologies). The 5' PCR primer contained a recognition sequence for the restriction

endonuclease BamH1 5' to the aurora2 coding sequence. This allowed the insertion of the aurora2 gene in frame with the 6 histidine residues, spacer region and rTEV protease cleavage site encoded by the pFastBac HTc vector. The 3' PCR primer replaced the aurora2 stop codon with additional coding sequence followed by a stop codon and a recognition sequence for the restriction endonuclease Not1 . This additional coding sequence (5' TAC CCA TAC GAT

GTT CCA GAT TAC GCT TCT TAA 3' ) encoded for the polypeptide sequence YPYDVDPDYAS. This sequence, derived from the influenza hemagglutinin protein, is frequently used as a tag epitope sequence that can be identified using specific monoclonal antibodies. The recombinant pFastBac vector therefore encoded for an N-terminally 6 his tagged, C terminally influenza hemagglutinin epitope tagged aurora2 protein. Details of the methods for the assembly of recombinant DNA molecules can be found in standard texts, for example Sambrook et al. 1989, Molecular Cloning - A Laboratory Manual, 2<sup>nd</sup> Edition, Cold Spring Harbor Laboratory press and Ausubel et al. 1999, Current Protocols in Molecular Biology, John Wiley and Sons Inc.

Production of recombinant virus can be performed following manufacturer's protocol from GibcoBRL. Briefly, the pFastBac-1 vector carrying the aurora2 gene was transformed into E. coli DH10Bac cells containing the baculovirus genome (bacmid DNA) and via a transposition event in the cells, a region of the pFastBac vector containing gentamycin resistance gene and the aurora2 gene including the baculovirus polyhedrin promoter was transposed directly into the bacmid DNA. By selection on gentamycin, kanamycin, tetracycline and X-gal, resultant white colonies should contain recombinant bacmid DNA encoding aurora2. Bacmid DNA was extracted from a small scale culture of several BH10Bac white colonies and transfected into Spodoptera frugiperda Sf21 cells grown in TC100 medium (GibcoBRL) containing 10% serum using CellFECTIN reagent (GibcoBRL) following manufacturer's instructions. Virus particles were harvested by collecting cell culture medium 72 hrs post transfection. 0.5 mls of medium was used to infect 100 ml suspension culture of Sf21s containing  $1 \times 10^7$  cells/ml. Cell culture medium was harvested 48 hrs post infection and virus titre determined using a standard plaque assay procedure. Virus stocks were used to infect Sf9 and "High 5" cells at a multiplicity of infection (MOI) of 3 to ascertain expression of recombinant aurora2 protein.

For the large scale expression of aurora2 kinase activity, Sf21 insect cells were grown at 28°C in TC100 medium supplemented with 10% foetal calf serum (Viralex) and 0.2% F68

Pluronic (Sigma) on a Wheaton roller rig at 3 r.p.m. When the cell density reached  $1.2 \times 10^6$  cells  $\text{ml}^{-1}$  they were infected with plaque-pure aurora2 recombinant virus at a multiplicity of infection of 1 and harvested 48 hours later. All subsequent purification steps were performed at  $4^\circ\text{C}$ . Frozen insect cell pellets containing a total of  $2.0 \times 10^8$  cells were thawed and diluted

5 with lysis buffer (25 mM HEPES (N-[2-hydroxyethyl]piperazine-N'-[2-ethanesulphonic acid]) pH7.4 at  $4^\circ\text{C}$ , 100 mM KCl, 25 mM NaF, 1 mM  $\text{Na}_3\text{VO}_4$ , 1 mM PMSF (phenylmethylsulphonyl fluoride), 2 mM 2-mercaptoethanol, 2 mM imidazole, 1  $\mu\text{g}/\text{ml}$  aprotinin, 1  $\mu\text{g}/\text{ml}$  pepstatin, 1  $\mu\text{g}/\text{ml}$  leupeptin), using 1.0 ml per  $3 \times 10^7$  cells. Lysis was achieved using a dounce homogeniser, following which the lysate was centrifuged at 41,000g

10 for 35 minutes. Aspirated supernatant was pumped onto a 5 mm diameter chromatography column containing 500  $\mu\text{l}$  Ni NTA (nitrilo-tri-acetic acid) agarose (Qiagen, product no. 30250) which had been equilibrated in lysis buffer. A baseline level of UV absorbance for the eluent was reached after washing the column with 12 ml of lysis buffer followed by 7 ml of wash buffer (25 mM HEPES pH7.4 at  $4^\circ\text{C}$ , 100 mM KCl, 20 mM imidazole, 2 mM 2-

15 mercaptoethanol). Bound aurora2 protein was eluted from the column using elution buffer (25 mM HEPES pH7.4 at  $4^\circ\text{C}$ , 100 mM KCl, 400 mM imidazole, 2 mM 2-mercaptoethanol). An elution fraction (2.5 ml) corresponding to the peak in UV absorbance was collected. The elution fraction, containing active aurora2 kinase, was dialysed exhaustively against dialysis buffer (25 mM HEPES pH7.4 at  $4^\circ\text{C}$ , 45% glycerol (v/v), 100 mM KCl, 0.25% Nonidet P40

20 (v/v), 1 mM dithiothreitol).

Each new batch of aurora2 enzyme was titrated in the assay by dilution with enzyme diluent (25mM Tris-HCl pH7.5, 12.5mM KCl, 0.6mM DTT). For a typical batch, stock enzyme is diluted 1 in 666 with enzyme diluent & 20 $\mu\text{l}$  of dilute enzyme is used for each assay well. Test compounds (at 10mM in dimethylsulphoxide (DMSO)) were diluted with

25 water & 10 $\mu\text{l}$  of diluted compound was transferred to wells in the assay plates. "Total" & "blank" control wells contained 2.5% DMSO instead of compound. Twenty microlitres of freshly diluted enzyme was added to all wells, apart from "blank" wells. Twenty microlitres of enzyme diluent was added to "blank" wells. Twenty microlitres of reaction mix (25mM Tris-HCl, 78.4mM KCl, 2.5mM NaF, 0.6mM dithiothreitol, 6.25mM  $\text{MnCl}_2$ , 6.25mM ATP,

30 7.5 $\mu\text{M}$  peptide substrate [biotin-LRRWSLGLRRWSLGLRRWSLGLRRWSLG]) containing 0.2 $\mu\text{Ci}$  [ $\gamma^{33}\text{P}$ ]ATP (Amersham Pharmacia, specific activity  $\geq 2500\text{Ci}/\text{mmol}$ ) was then added to

all test wells to start the reaction. The plates were incubated at room temperature for 60 minutes. To stop the reaction 100µl 20% v/v orthophosphoric acid was added to all wells. The peptide substrate was captured on positively-charged nitrocellulose P30 filtermat (Whatman) using a 96-well plate harvester (TomTek) & then assayed for incorporation of  $^{33}\text{P}$  with a Beta plate counter. "Blank" (no enzyme) and "total" (no compound) control values were used to determine the dilution range of test compound which gave 50% inhibition of enzyme activity.

In this test, compound 4 in Table 1 gave 50% inhibition of enzyme activity at a concentration of 0.465µM

(b) In Vitro cell proliferation assay

This assay determines the ability of a test compound to inhibit the growth of adherent mammalian cell lines, for example the human tumour cell line MCF7.

MCF-7 (ATCC HTB-22) or other adherent cells were typically seeded at  $1 \times 10^3$  cells per well (excluding the peripheral wells) in DMEM (Sigma Aldrich) without phenol red, plus 10% foetal calf serum, 1% L-glutamine and 1% penicillin/streptomycin in 96 well tissue culture treated clear plates (Costar). The following day (day 1), the media was removed from a no treatment control plate and the plate stored at -80°C. The remaining plates were dosed with compound (diluted from 10mM stock in DMSO using DMEM (without phenol red, 10% FCS, 1% L-glutamine, 1% penicillin/streptomycin). Untreated control wells were included on each plate. After 3 days in the presence / absence of compound (day 4) the media was removed and the plates stored at -80°C. Twenty four hours later the plates were thawed at room temperature and cell density determined using the CyQUANT cell proliferation assay kit (c-7026/c-7027 Molecular Probes Inc.) according to manufacturers directions. Briefly, 200µl of a cell lysis / dye mixture (10µl of 20X cell lysis buffer B, 190µl of sterile water, 0.25µl of CYQUANT GR dye) was added to each well and the plates incubated at room temperature for 5 minutes in the dark. The fluorescence of the wells was then measured using a fluorescence microplate reader (gain 70, 2 reads per well, 1 cycle with excitation 485nm and emission 530nm using a CytoFluor plate reader (PerSeptive Biosystems Inc.)). The values from day 1 and day 4 (compound treated) together with the values from the untreated cells were used to determine the dilution range of a test compound that gave 50% inhibition of cell proliferation. Compound no.4 in Table 1 was effective in this test at 12.4µM



These values could also be used to calculate the dilution range of a test compound at which the cell density dropped below the day 1 control value. This indicates the cytotoxicity of the compound.

5    (c) In Vitro cell cycle analysis assay

---

This assay determines the ability of a test compound to arrest cells in specific phases of the cell cycle. Many different mammalian cell lines could be used in this assay and MCF7 cells are included here as an example. MCF-7 cells were seeded at  $3 \times 10^5$  cells per T25 flask (Costar) in 5 ml DMEM (no phenol red 10% FCS, 1% L-glutamine 1% penicillin / streptomycin). Flasks were then incubated overnight in a humidified 37°C incubator with 5% CO<sub>2</sub>. The following day 1ml of DMEM (no phenol red 10% FCS, 1% L-glutamine 1% penicillin / streptomycin) carrying the appropriate concentration of test compound solubilised in DMSO was added to the flask. A no compound control treatments was also included (0.5% DMSO). The cells were then incubated for a defined time (usually 24 hours) with compound. After this time the media was aspirated from the cells and they were washed with 5ml of prewarmed (37°C) sterile PBSA, then detached from the flask by brief incubation with trypsin and followed by resuspension in 10ml of 1% Bovine Serum Albumin (BSA, Sigma-Aldrich Co.) in sterile PBSA. The samples were then centrifuged at 2200rpm for 10 min. The supernatant was aspirated and the cell pellet was resuspended in 200µl of 0.1% (w/v) Tris sodium citrate, 0.0564% (w/v) NaCl, 0.03% (v/v) Nonidet NP40, [pH 7.6]. Propidium Iodide (Sigma Aldrich Co.) was added to 40µg/ml and RNAase A (Sigma Aldrich Co.) to 100µg/ml. The cells were then incubated at 37°C for 30 minutes. The samples were centrifuged at 2200rpm for 10 min, the supernatant removed and the remaining pellet (nuclei) resuspended in 200µl of sterile PBSA. Each sample was then syringed 10 times using 21 gauge needle. The samples were then transferred to LPS tubes and DNA content per cell analysed by Fluorescence activated cell sorting (FACS) using a FACScan flow cytometer (Becton Dickinson). Typically 25000 events were counted and recorded using CellQuest v1.1 software (Verity Software). Cell cycle distribution of the population was calculated using Modfit software (Verity Software) and expressed as percentage of cells in G0/G1, S and G2/M phases of the cell cycle.

Treating MCF7 cells with 24.8 $\mu$ M Compound no.4 in Table 1 for 24 hours produced the following changes in cell cycle distribution:

Treatment	% Cells in G1	% Cells in S	% Cells in G2/M
DMSO (control)	60.96	26.99	12.05
24.8 $\mu$ M Compound 4	37.29	33.93	28.78